

CYCLOTOMIC BIRMAN–WENZL–MURAKAMI ALGEBRAS, II: ADMISSIBILITY RELATIONS AND FREENESS

FREDERICK M. GOODMAN AND HOLLY HAUSCHILD MOSLEY

ABSTRACT. The cyclotomic Birman–Wenzl–Murakami algebras are quotients of the affine BMW algebras in which the affine generator satisfies a polynomial relation. We study admissibility conditions on the ground ring for these algebras, and show that the algebras defined over an admissible integral ground ring S are free S –modules and isomorphic to cyclotomic Kauffman tangle algebras. We also determine the representation theory in the generic semisimple case, obtain a recursive formula for the weights of the Markov trace, and give a sufficient condition for semisimpliity.

March, 2008.

CONTENTS

1. Introduction	1
2. Preliminaries	6
3. Representations of the 2–strand algebra and admissibility conditions	16
4. The generic structure of the cyclotomic BMW algebra	32
5. The $\mathbb{Z}_r$ –Brauer algebra and the dimension of $W_{n,F,r}$.	42
6. Freeness and isomorphism with cyclotomic Kauffman tangle algebras.	45
References	45

1. INTRODUCTION

This paper and the companion paper [10] continue the study of affine and cyclotomic Birman–Wenzl– Murakami (BMW) algebras, which we began in [9].

1.1. Background. The origin of the BMW algebras was in knot theory. Kauffman defined [14] an invariant of regular isotopy for links in S^3 , determined by certain skein relations. Birman and Wenzl [6] and independently Murakami [22] then defined a family of quotients of the braid group algebras, and showed that

Kauffman’s invariant could be recovered from a trace on these algebras. These (BMW) algebras were defined by generators and relations, but were implicitly modeled on certain algebras of tangles whose definition was subsequently made explicit by Morton and Traczyk [20], as follows: Let S be a commutative unital ring with invertible elements ρ , q , and δ_0 satisfying $\rho^{-1} - \rho = (q^{-1} - q)(\delta_0 - 1)$. The *Kauffman tangle algebra* $KT_{n,S}$ is the S -algebra of framed (n, n) -tangles in the disc cross the interval, modulo Kauffman skein relations:

- (1) Crossing relation: $\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} - \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array} = (q^{-1} - q) \left(\begin{array}{c} \diagup \diagup \\ \diagdown \diagdown \end{array} - \begin{array}{c} \diagdown \diagdown \\ \diagup \diagup \end{array} \right).$
- (2) Untwisting relation: $\begin{array}{c} \text{positive twist} \end{array} = \rho \quad \Bigg| \quad \text{and} \quad \begin{array}{c} \text{negative twist} \end{array} = \rho^{-1} \quad \Bigg|.$
- (3) Free loop relation: $T \cup \bigcirc = \delta_0 T.$

Morton and Traczyk [20] showed that the n -strand algebra $KT_{n,S}$ is free of rank $(2n - 1)!!$ as a module over S , and Morton and Wassermann [21] proved that the BMW algebras and the Kauffman tangle algebras are isomorphic.

It is natural to “affinize” the BMW algebras to obtain BMW analogues of the affine Hecke algebras of type A , see [1]. The affine Hecke algebra can be realized geometrically as the algebra of braids in the annulus cross the interval, modulo Hecke skein relations; this suggests defining the affine Kauffman tangle algebra $\widehat{KT}_{n,S}$ as the algebra of framed (n, n) -tangles in the annulus cross the interval, modulo Kauffman skein relations. Turaev [28] showed that the resulting algebra of $(0, 0)$ -tangles is a (commutative) polynomial algebra in infinitely many variables, so it makes sense to absorb this polynomial algebra into the ground ring. (The ground ring gains infinitely many parameters δ_j ($j \geq 1$) corresponding to the generators of the polynomial algebra.) On the other hand, Häring–Oldenburg [12] defined an affine version of the BMW algebras by generators and relations. In [9], we showed that Häring–Oldenburg’s affine BMW algebras are isomorphic to the affine Kauffman tangle algebras, and we showed that these algebras are free modules over their ground ring, with a basis reminiscent of a well-known basis of affine Hecke algebras.

The affine BMW algebras arise naturally in several different contexts: knot theory in the solid torus [28, 16, 17], representations of Artin braid group of type B by R matrices of quantum groups [24], and representations of ordinary BMW algebras. See [10], Section 1.1 for more detail.

1.2. Cyclotomic algebras. In this paper and the companion paper [10] we consider cyclotomic BMW algebras, which are the BMW analogues of cyclotomic Hecke algebras [1]. The affine BMW algebras have a distinguished generator x_1 , which, in the geometric (Kauffman tangle) picture is represented by a braid with one strand wrapping around the hole in the annulus cross interval. The *cyclotomic BMW algebra* $W_{n,S,r}$ is defined to be the quotient of the affine BMW algebra $\widehat{W}_{n,S}$ in which the generator x_1 satisfies a monic polynomial equation

$$(1.1) \quad x_1^r + \sum_{k=0}^{r-1} a_k x_1^k = 0.$$

with coefficients in S .¹ The cyclotomic BMW algebras were also introduced by Häring-Oldenburg in [12].

In the geometric (Kauffman tangle) picture, it is more natural to convert this relation into a local skein relation:

$$(1.2) \quad T_r + \sum_{k=0}^{r-1} a_k T_k = 0,$$

whenever $T_0, T_1, \dots, T_r$ are affine tangle diagrams that are identical in the exterior of some disc E and $T_k \cap E$ consists of one strand wrapping k times around the hole in the annulus cross interval; i.e. $T_k \cap E$ “equals” x_1^k . The *cyclotomic Kauffman tangle algebra* $KT_{n,S,r}$ is defined to be the quotient of the affine Kauffman tangle algebra $\widehat{KT}_{n,S}$ by the cyclotomic skein relation. See [10], Section 1.2, for an alternative description of the cyclotomic Kauffman tangle algebras in terms of the affine Kauffman tangle category.

A priori, the ideal in $\widehat{KT}_{n,S} \cong \widehat{W}_{n,S}$ generated by the cyclotomic skein relation (1.2) is larger than the ideal generated by the polynomial relation (1.1), so we have a surjective, but not evidently injective homomorphism

$$\varphi : W_{n,S,r} \rightarrow KT_{n,S,r}.$$

1.3. Admissibility. The cyclotomic BMW algebras and Kauffman tangle algebras can be defined over an arbitrary commutative unital ring S with parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$ (the eigenvalues of y_1), assuming that ρ, q, δ_0 and $u_1, \dots, u_r$ are invertible, and $\rho^{-1} - \rho = (q^{-1} - q)(\delta_0 - 1)$. However, unless the parameters satisfy additional relations, the identity element $\mathbf{1}$ of the cyclotomic Kauffman tangle algebras will be a torsion element over S ; if S is a field (and

¹Actually, we will assume that the polynomial splits in S .

the additional relations do not hold) then $\mathbf{1} = 0$, so $KT_{n,S,r}$ becomes trivial. The additional conditions are called “weak admissibility;” see Section 2.5. Weak admissibility is thus a minimal condition for the non-triviality of the cyclotomic algebras.

In order to obtain substantial results about the cyclotomic BMW and Kauffman tangle algebras, it seems necessary to impose a condition on the ground ring S that is stronger than weak admissibility. An appropriate condition was introduced by Wilcox and Yu in [31]. Their condition has a simple formulation in terms of the representation theory of the 2-strand algebra $W_{2,S,r}$, and also translates into explicit relations on the parameters. The condition of Wilcox and Yu, called “admissibility,” is the subject of Section 3 of this paper.

1.4. Results. The main results of this paper and [10] is that if the ground ring S is an integral domain and admissible in the sense of Wilcox and Yu, then $W_{n,S,r} \cong KT_{n,S,r}$, and, moreover, these algebras are free S -modules of rank $r^n(2n-1)!!$. The proof of these results has a topological component, which is given in [10], and an algebraic component, which is given in this paper.

The topological arguments in [10] have two essential consequences: first they show that the cyclotomic BMW algebra $W_{n,S,r}$ has a spanning set $\mathbb{A}'_r$ of cardinality $r^n(2n-1)!!$. Second, they show that the $(0,0)$ -tangle algebra $KT_{0,S,r}$ is free of rank 1, assuming that S is weakly admissible. This implies the existence of the “Markov trace” on the cyclotomic BMW and Kauffman tangle algebras (and, hence, Kauffman type invariants for links in the solid torus).

To achieve the main results, it remains to show that the spanning set $\varphi(\mathbb{A}'_r)$ of the cyclotomic Kauffman tangle algebra $KT_{n,S,r}$ is linearly independent, when the ground ring S is an admissible integral domain.

To do this we first show that there exists a universal admissible integral domain $\overline{R}_+$ with the property that every admissible integral domain is a quotient of $\overline{R}_+$. Moreover, field of fractions F of $\overline{R}_+$ is the field of rational functions $\mathbb{Q}(\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r)$.

We then analyze the representation theory of the cyclotomic BMW algebras defined over F , by adapting the inductive method of Wenzl from [6, 29, 30]. This analysis shows that $KT_{n,F,r} \cong W_{n,F,r}$, and that these algebras are split semisimple of dimension $r^n(2n-1)!!$. The irreducible representations of $W_{n,F,r}$ are indexed by r -tuples of Young diagrams of total size $\leq n$ and congruent to $n \bmod 2$.

A relatively simple argument then shows that for any admissible integral domain S , we have $KT_{n,S,r} \cong W_{n,S,r}$, and these algebras are free of rank $r^n(2n-1)!!$ over S .

As a biproduct of our argument, we obtain a sufficient condition for semisimplicity of the cyclotomic BMW algebras defined over an admissible field.

The elements

$$y_j = g_{j-1} \cdots g_1 y_1 g_1 \cdots g_{j-1},$$

are important for the representation theory of the cyclotomic BMW algebras. They are BMW analogues of Jucys-Murphy elements; see, for example, [18, 19]. Indeed, in our treatment of the representation theory over the generic field F , the action of the Jucys-Murphy elements plays an essential role.

1.5. Related work, and acknowledgments. Wilcox and Yu have been studying the same material independently and have obtained similar (and, in some instances, slightly stronger) results [31, 33, 32]. We are indebted to Wilcox and Yu for pointing out an error in a previous preprint version of our work, which required us to substantially rework the algebraic (linear independence) component of the arguments. In fact, we had to adopt a completely different strategy for proving linear independence. Meanwhile, Wilcox and Yu [33, 32] were able to make our original strategy work, using a refined analysis of their admissibility condition.

We would also like to mention here recent work of Ariki, Mathas and Rui on the “cyclotomic Nazarov–Wenzl algebras” [3], which are cyclotomic quotients of the degenerate affine BMW algebras introduced by Nazarov [23]. The idea of deriving an admissibility condition on the ground ring from an assumption on the representation theory of the 2-strand algebra originates in [3], and is essential to our work and the work of Wilcox and Yu.

We rely heavily on ideas from Beliakova and Blanchet [5] (and indirectly on Nazarov [23]) for our computation of the weights of the Markov trace and of the action of conjugates of the affine generator y_1 on the basis of up–down tableaux.

It is shown in [7] and [33] that cyclotomic BMW algebras defined over integral admissible ground rings are cellular in the sense of Graham and Lehrer [11]. The proof in [7] depends on the isomorphism of cyclotomic BMW algebras and cyclotomic Kauffmann tangle algebras established here.

Finally, a recent preprint of Rui and Xu [26] concerns the representation theory of cyclotomic BMW algebras over a u -admissible field.

2. PRELIMINARIES

2.1. The affine BMW algebra. We recall the definition of the affine Birman–Wenzl–Murakami (BMW) algebra. Let S be a commutative unital ring with elements ρ , q , and δ_j ($j \geq 0$), such that ρ , q , and δ_0 are invertible, and the relation $\rho^{-1} - \rho = (q^{-1} - q)(\delta_0 - 1)$ holds.

Definition 2.1. The *affine Birman–Wenzl–Murakami algebra* $\widehat{W}_{n,S}$ is the S algebra with generators $y_1^{\pm 1}$, $g_i^{\pm 1}$ and e_i ($1 \leq i \leq n-1$) and relations:

- (1) (Inverses) $g_i g_i^{-1} = g_i^{-1} g_i = 1$ and $y_1 y_1^{-1} = y_1^{-1} y_1 = 1$.
- (2) (Idempotent relation) $e_i^2 = \delta_0 e_i$.
- (3) (Affine braid relations)
 - (a) $g_i g_{i+1} g_i = g_{i+1} g_i g_{i+1}$ and $g_i g_j = g_j g_i$ if $|i - j| \geq 2$.
 - (b) $y_1 g_1 y_1 g_1 = g_1 y_1 g_1 y_1$ and $y_1 g_j = g_j y_1$ if $j \geq 2$.
- (4) (Commutation relations)
 - (a) $g_i e_j = e_j g_i$ and $e_i e_j = e_j e_i$ if $|i - j| \geq 2$.
 - (b) $y_1 e_j = e_j y_1$ if $j \geq 2$.
- (5) (Affine tangle relations)
 - (a) $e_i e_{i \pm 1} e_i = e_i$,
 - (b) $g_i g_{i \pm 1} e_i = e_{i \pm 1} e_i$ and $e_i g_{i \pm 1} g_i = e_i e_{i \pm 1}$.
 - (c) For $j \geq 1$, $e_1 y_1^j e_1 = \delta_j e_1$.
- (6) (Kauffman skein relation) $g_i - g_i^{-1} = (q - q^{-1})(1 - e_i)$.
- (7) (Untwisting relations) $g_i e_i = e_i g_i = \rho^{-1} e_i$ and $e_i g_{i \pm 1} e_i = \rho e_i$.
- (8) (Unwrapping relation) $e_1 y_1 g_1 y_1 = \rho e_1 = y_1 g_1 y_1 e_1$.

Remark 2.2. The presentation differs slightly from the one we used in [9]. There we used the generator $x_1 = \rho^{-1} y_1$, and parameters $\vartheta_j = \rho^{-j} \delta_j$ (so that $e_1 x_1^j e_1 = \vartheta_j e_1$). We also used the parameter z in place of $(q^{-1} - q)$.

In $\widehat{W}_{n,S}$, define for $1 \leq j \leq n$,

$$y_j = g_{j-1} \cdots g_1 y_1 g_1 \cdots g_{j-1}$$

and

$$y'_j = g_{j-1} \cdots g_1 y_1 g_1^{-1} \cdots g_{j-1}^{-1}.$$

We record some elementary results from [9]:

Lemma 2.3. *The following relations hold in $\widehat{W}_{n,S}$. For $1 \leq k, j \leq n$,*

- (1) *For $j \notin \{k, k-1\}$, $g_j y_k = y_k g_j$.*
- (2) *For $j \notin \{k, k-1\}$, $e_j y_k = y_k e_j$.*
- (3) *$y_j y_k = y_k y_j$.*
- (4) *For $k < n$, $g_k y_k = y_{k+1} g_k^{-1}$ and $g_k^{-1} y_{k+1} = y_k g_k$.*
- (5) *For $k < n$,*

$$\begin{aligned} g_k y_{k+1} &= y_k g_k + (q - q^{-1}) y_{k+1} - (q - q^{-1}) \rho e_k y_{k+1}, \\ g_k^{-1} y_k &= y_{k+1} g_k^{-1} - (q - q^{-1}) y_k + (q - q^{-1}) e_k y_k. \end{aligned}$$

- (6) *For $k < n$, $e_k y_k = e_k y_{k+1}^{-1}$ and $e_k y_k^{-1} = e_k y_{k+1}$.*
- (7) *$g_k y_k g_k y_k = y_k g_k y_k g_k$.*
- (8) *$e_k y_k g_k y_k = \rho e_k$.*
- (9) *For $s \geq 1$, $e_k y_k^s g_k y_k = e_k y_k^{s-1} g_k^{-1}$, and $e_k y_k^{-s} g_k^{-1} y_k^{-1} = e_k y_k^{-s+1} g_k$.*

Remark 2.4. There is a unique antiautomorphism of the affine BMW algebra leaving each of the generators y_1, e_i, g_i invariant; the antiautomorphism simply reverses the order of a word in the generators. This follows from the symmetry of the relations. Using this, one obtains versions of points (6), (8) and (9) of the Lemma with e_k written on the right, for example: $y_k g_k y_k^s e_k = g_k^{-1} y_k^{s-1} e_k$.

Lemma 2.5. *For $j \geq 1$, there exist elements $\delta_{-j} \in \mathbb{Z}[\rho^{\pm 1}, q^{\pm 1}, \delta_0, \dots, \delta_j]$ such that $e_1 y_1^{-j} e_1 = \delta_{-j} e_1$. Moreover, the elements δ_{-j} are determined by the recursion relation:*


$$(2.1) \quad \begin{aligned} \delta_{-1} &= \rho^{-2} \delta_1 \\ \delta_{-j} &= \rho^{-2} \delta_j + (q^{-1} - q) \rho^{-1} \sum_{k=1}^{j-1} (\delta_k \delta_{k-j} - \delta_{2k-j}) \quad (j \geq 2). \end{aligned}$$

Proof. Follows from [9], Corollary 3.13, and [10], Lemma 2.6. $\square$

2.2. The affine Kauffman tangle algebra. Let S be a commutative unital ring with elements ρ, q , and δ_j ($j \geq 0$), such that ρ, q , and δ_0 are invertible, and the relation $\rho^{-1} - \rho = (q^{-1} - q)(\delta_0 - 1)$ holds. The *affine Kauffman tangle algebra* $\widehat{KT}_{n,S}$ is the S -algebra of framed (n, n) tangles in $A \times I$, where A is the annulus and I the interval, modulo the Kauffman skein relations. The reader is referred to [9] for a detailed definition.

The affine Kauffman tangle algebra $\widehat{KT}_{n,S}$ can be described in terms of *affine tangle diagrams*, which are quasi-planar diagrams representing framed tangles.

Here is a picture of a typical affine tangle diagram. The heavy vertical line represents the hole in $A \times I$; we refer to it as the flagpole.



Multiplication is by stacking of such diagrams; our convention is that the product ab is given by stacking b over a . The affine Kauffman tangle algebra is generated by the following affine tangle diagrams:

$$X_1 = \begin{array}{c} | \\ \text{---} \text{---} \text{---} \end{array} \quad G_i = \begin{array}{c} | \quad | \quad | \quad | \\ \text{---} \text{---} \text{---} \end{array} \quad E_i = \begin{array}{c} | \quad | \quad | \quad | \\ \text{---} \text{---} \end{array}$$

$i \quad i+1 \qquad i \quad i+1$

Theorem 2.6 ([9]). *The affine BMW algebra $\widehat{W}_{n,S}$ is isomorphic to the affine Kauffman tangle algebra $\widehat{KT}_{n,S}$ by a map determined by the assignments $y_1 \mapsto \rho X_1$, $e_i \mapsto E_i$, and $g_i \mapsto G_i$ for $1 \leq i \leq n-1$.*

2.3. The cyclotomic BMW and Kauffman tangle algebras. We will consider algebras defined over ground rings with parameters satisfying certain conditions. In order to avoid repeating these conditions, we establish the following convention:

Definition 2.7. A *ground ring* S is a commutative unital ring with parameters ρ , q , δ_j ($j \geq 0$), and $u_1, \dots, u_r$, with ρ , q , δ_0 , and $u_1, \dots, u_r$ invertible, and with $\rho^{-1} - \rho = (q^{-1} - q)(\delta_0 - 1)$.

Definition 2.8. Let S be a ground ring with parameters ρ , q , δ_j ($j \geq 0$), and $u_1, \dots, u_r$. The *cyclotomic BMW algebra* $W_{n,S,r}(u_1, \dots, u_r)$ is the quotient of $\widehat{W}_{n,S}$ by the relation

$$(2.2) \quad (y_1 - u_1)(y_1 - u_2) \cdots (y_1 - u_r) = 0.$$

Remark 2.9. The assignment $e_i \mapsto e_i$, $g_i \mapsto g_i$, $y_1 \mapsto y_1$ defines a homomorphism ι from $W_{n,S,r}$ to $W_{n+1,S,r}$, since the relations are preserved. It is not evident that ι is injective. However, when S is an admissible integral domain, we will show that $W_{n,S,r}$ is isomorphic to a cyclotomic version of the Kauffman tangle algebra (defined below), and in this case, it is true that ι is injective.

Remark 2.10. Let S be a ring with parameters ρ, q , etc., as above, and let S' be another ring with parameters ρ', q' , etc. Suppose there is a ring homomorphism $\psi : S \rightarrow S'$ mapping $\rho \mapsto \rho', q \mapsto q'$, etc. Then $W_{n,S',r}$ can be regarded as an algebra over S , with $sx = \psi(s)x$ for $s \in S$ and $x \in W_{n,S',r}$, and there is an S -algebra homomorphism $\tilde{\psi} : W_{n,S,r} \rightarrow W_{n,S',r}$ taking generators to generators. We claim that $W_{n,S,r} \otimes_S S' \cong W_{n,S',r}$ as S' -algebras. In fact, we have the S' algebra homomorphism $\tilde{\psi} \otimes \text{id} : W_{n,S,r} \otimes_S S' \rightarrow W_{n,S',r} \otimes_S S' \cong W_{n,S',r}$. In the other direction, we have an S' -algebra homomorphism $\theta : W_{n,S',r} \rightarrow W_{n,S,r} \otimes_S S'$ mapping $g_i \mapsto g_i \otimes 1$, etc. The maps $\tilde{\psi} \otimes \text{id}$ and θ are inverses.

Of course, this remark applies in general to algebras defined by generators and relations.²

Now we consider how to define a cyclotomic version of the Kauffman tangle algebra. Rewrite the relation (2.2) in the form

$$\sum_{k=0}^r (-1)^{r-k} \varepsilon_{r-k}(u_1, \dots, u_r) y_1^k = 0,$$

where ε_j is the j -th elementary symmetric function. The corresponding relation in the affine Kauffman tangle algebra is

$$(2.3) \quad \sum_{k=0}^r (-1)^{r-k} \varepsilon_{r-k}(u_1, \dots, u_r) \rho^k X_1^k = 0,$$

Now we want to impose this as a local skein relation.

Definition 2.11. Let S be a ground ring with parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. The *cyclotomic Kauffman tangle algebra* $KT_{n,S,r}(u_1, \dots, u_r)$ is the quotient of the affine Kauffman tangle algebra $\widehat{KT}_{n,S}$ by the cyclotomic skein relation:

$$(2.4) \quad \sum_{k=0}^r (-1)^{r-k} \varepsilon_{r-k}(u_1, \dots, u_r) \rho^k \left(\text{disc with } X_1^k \right) = 0,$$

The sum is over affine tangle diagrams which differ only in the interior of the indicated disc and are identical outside of the disc; the interior of the disc contains an interval on the flagpole and a piece of an affine tangle diagram isotopic to X_1^k .

² We could have slightly simplified some arguments in [9] using this remark.

We continue to write E_i, G_i, X_1 for the image of these elements of the affine Kauffman tangle algebra in the cyclotomic Kauffman tangle algebra. We write

$$Y_1 = \rho X_1.$$

The ideal in the affine Kauffman tangle algebra by which we are taking the quotient contains the ideal generated by $(Y_1 - u_1)(Y_1 - u_2) \cdots (Y_1 - u_r)$, but could in principal be larger. It follows that there is a homomorphism

$$\varphi : W_{n,S,r}(u_1, \dots, u_r) \rightarrow KT_{n,S,r}(u_1, \dots, u_r)$$

determined by $y_1 \mapsto Y_1, e_i \mapsto E_i, g_i \mapsto G_i$. Moreover, the following diagram (in which the vertical arrows are the quotient maps) commutes.

$$\begin{array}{ccc} \widehat{W}_{n,S} & \xrightarrow{\varphi} & \widehat{KT}_{n,S} \\ \downarrow \pi & & \downarrow \pi \\ W_{n,S,r}(u_1, \dots, u_r) & \xrightarrow{\varphi} & KT_{n,S,r}(u_1, \dots, u_r) \end{array}$$

The homomorphism $\varphi : W_{n,S,r}(u_1, \dots, u_r) \rightarrow KT_{n,S,r}(u_1, \dots, u_r)$ is surjective because the diagram commutes and $\varphi : \widehat{W}_{n,S} \rightarrow \widehat{KT}_{n,S}$ is an isomorphism.

Remark 2.12. Let S be a ring with parameters ρ, q , etc., as above, and let S' be another ring with parameters ρ', q' , etc. Suppose there is a ring homomorphism $\psi : S \rightarrow S'$ mapping $\rho \mapsto \rho', q \mapsto q'$, etc. Any S' -algebra can be regarded as an S -algebra using ψ . We have a map of monoid rings $\tilde{\psi} : S \widehat{\mathcal{U}}(n, n) \rightarrow S' \widehat{\mathcal{U}}(n, n)$, and this map respects regular isotopy, the Kauffman skein relations, and the cyclotomic relations, so induces an S -algebra homomorphism from $\tilde{\psi} : KT_{n,S,r} \rightarrow KT_{n,S',r}$. As in Remark 2.10, we have $KT_{n,S,r} \otimes_S S' \cong KT_{n,S',r}$ as S' -algebras.

2.4. Morphisms of ground rings. We consider what are the appropriate morphisms between ground rings for cyclotomic BMW or Kauffman tangle algebras. The obvious notion would be that of a ring homomorphism taking parameters to parameters; that is, if S is a ring with parameters ρ, q , etc., and S' a weakly admissible ring with parameters ρ', q' , etc., then a morphism $\varphi : S \rightarrow S'$ would be required to map $\rho \mapsto \rho', q \mapsto q'$, etc.

However, it is better to require less, for the following reason: The parameter q enters into the cyclotomic BMW or Kauffman tangle relations only in the

expression $q^{-1} - q$, and the transformation $q \mapsto -q^{-1}$ leaves this expression invariant. Moreover, the transformation $g_i \mapsto -g_i$, $\rho \mapsto -\rho$, $q \mapsto -q$ (with all other generators and parameters unchanged) leaves the cyclotomic BMW relations unchanged. Likewise, for the cyclotomic Kauffman tangle algebras, the map taking an affine tangle diagram T to $(-1)^{\kappa(T)}T$, where $\kappa(T)$ is the number of crossings of ordinary strands of T , determines an isomorphism $KT_{n,S,r}(\rho, q, \dots) \cong KT_{n,S,r}(-\rho, -q, \dots)$.

Taking this into account, we arrive at the following notion:

Definition 2.13. Let S be a ground ring with parameters ρ , q , δ_j ($j \geq 0$), and $u_1, \dots, u_r$. Let S' be another ground ring with parameters ρ' , q' , etc.

A unital ring homomorphism $\varphi : S \rightarrow S'$ is a *morphism of ground rings* if it maps

$$\begin{cases} \rho \mapsto \rho', \text{ and} \\ q \mapsto q' \text{ or } q \mapsto -q'^{-1}, \end{cases}$$

or

$$\begin{cases} \rho \mapsto -\rho', \text{ and} \\ q \mapsto -q' \text{ or } q \mapsto q'^{-1}, \end{cases}$$

and strictly preserves all other parameters.

Suppose there is a morphism of ground rings $\psi : S \rightarrow S'$. Then ψ extends to a homomorphism from $W_{n,S,r}$ to $W_{n,S',r}$, and likewise to a homomorphism from $KT_{n,S,r}$ to $KT_{n,S',r}$. Moreover, $W_{n,S,r} \otimes_S S' \cong W_{n,S',r}$ and $KT_{n,S,r} \otimes_S S' \cong KT_{n,S',r}$ as S' -algebras. This follows from the observations above and Remarks 2.10 and 2.12.

2.5. Weak admissibility. As noted in [10], Section 4.1, unless the parameters of the ground ring S satisfy the *weak admissibility* condition defined below, the identity element $\mathbf{1}$ of the cyclotomic Kauffman tangle algebras will be a torsion element over S ; if S is a field (and the additional relations do not hold) then $\mathbf{1} = 0$, so $KT_{n,S,r} = \{0\}$. Similarly, in the absence of weak admissibility, a computation done in the cyclotomic BMW algebra shows that $e_1 \in W_{2,S,r}$ is a torsion element over S .


Definition 2.14. Let S be a ground ring with parameters ρ , q , δ_j , $j \geq 0$, and $u_1, \dots, u_r$. We say that the parameters are *weakly admissible* (or that the ring S

is weakly admissible) if the following relation holds:

$$\sum_{k=0}^r (-1)^{r-k} \varepsilon_{r-k}(u_1, \dots, u_r) \delta_{k+a} = 0,$$

for $a \in \mathbb{Z}$, where for $j \geq 1$, δ_{-j} is defined by the recursive relations of Lemma 2.5.

2.6. Inclusion, conditional expectation, and trace. Let S be a ground ring with parameters as above and write $KT_{n,S,r}$ for $KT_{n,S,r}(u_1, \dots, u_r)$. The affine Kauffman tangle algebras have inclusion maps $\iota : \widehat{KT}_{n-1} \rightarrow \widehat{KT}_n$, defined on the level of affine tangle diagrams by adding an additional strand on the right without adding any crossings:




Since these maps respect the cyclotomic relation (2.4), they induce homomorphisms

$$\iota : KT_{n-1,S,r} \rightarrow KT_{n,S,r}.$$

Recall also that the affine Kauffman tangle algebras have a conditional expectation $\varepsilon_n : \widehat{KT}_n \rightarrow \widehat{KT}_{n-1}$ defined by

$$\varepsilon_n(T) = \delta_0^{-1} \text{cl}_n(T),$$

where cl_n is the map of affine (n, n) -tangle diagrams to affine $(n-1, n-1)$ -tangle diagrams that “closes” the rightmost strand, without adding any crossings:



These maps respect the cyclotomic relation (2.4), so induce conditional expectations

$$\varepsilon_n : KT_{n,S,r} \rightarrow KT_{n-1,S,r}.$$

Since $\varepsilon_n \circ \iota$ is the identity on $KT_{n-1,S,r}$, it follows that $\iota : KT_{n-1,S,r} \rightarrow KT_{n,S,r}$ is injective.

It is shown in the companion paper [10], Proposition 4.3 that if the ground ring S is weakly admissible, then the cyclotomic Kauffman tangle algebra $KT_{0,S,r}$ is

a free S -module of rank 1. Define

$$\varepsilon = \varepsilon_1 \circ \cdots \circ \varepsilon_n : KT_{n,S,r} \rightarrow KT_{0,S,r} \cong S$$

It follows from [9], Proposition 2.14, that ε is a trace. We also define $\varepsilon : W_{n,S,r} \rightarrow S$ by $\varepsilon = \varepsilon \circ \varphi$, where $\varphi : W_{n,S,r} \rightarrow KT_{n,S,r}$ is the canonical homomorphism. Then ε is a trace on $W_{n,S,r}$ with the *Markov property*: for $b \in W_{n-1,S,r}$,

- (a) $\varepsilon(bg_{n-1}^{\pm 1}) = (\rho^{\pm 1}/\delta_0)\varepsilon(b)$,
- (b) $\varepsilon(be_{n-1}) = (1/\delta_0)\varepsilon(b)$,
- (c) $\varepsilon(b(y'_n)^r) = \delta_r\varepsilon(b)$, and

for $r \in \mathbb{Z}$, where $y'_n = (g_{n-1} \cdots g_1)y_1(g_1^{-1} \cdots g_{n-1}^{-1})$. See [9], Corollary 6.16.

Lemma 2.15. *Let S be a weakly admissible ring. For $x \in KT_{n,S,r}$,*

- (1) $E_n x E_n = \delta_0 \varepsilon_n(x)$.
- (2) $\varepsilon(E_n x E_n) = \varepsilon(x)$.
- (3) $\varepsilon(E_n x) = \delta_0^{-1} \varepsilon(x)$.

Proof. Straightforward. □

Lemma 2.16. *Let S be a weakly admissible ring. Then for all $n \geq 1$, E_n and G_n are non-zero elements in $KT_{n,S,r}$.*

Proof. $\varepsilon(E_n) = \delta_0^{-1}$ and $\varepsilon(G_n) = \rho^{-1} \delta_0^{-2}$. □

2.7. Inclusions of split semisimple algebras. A general source for the material in this section is [8]. The Jones basic construction (discussed below) is from [13].

A finite dimensional split semisimple algebra over a field F is one which is isomorphic to a finite direct sum of full matrix algebras over F . Suppose A and B are finite dimensional split semisimple algebras over F . Let A_i , $i \in I$, be the minimal ideals of A and B_j , $j \in J$, the minimal ideals of B . Let $\psi : A \rightarrow B$ be a unital homomorphism of algebras.

We associated a $J \times I$ *multiplicity matrix* Λ_ψ to ψ , as follows. Let W_j be a simple B_j -module. Then W_j becomes an A -module via ψ , and $\Lambda_\psi(j, i)$ is the multiplicity of a simple A_i -module in the decomposition of W_j as an A -module. In particular, if $A \subseteq B$ (and both algebras have the same identity element) the multiplicity matrix for the inclusion is called the *inclusion matrix* for $A \subseteq B$.

An equivalent characterization of the multiplicity matrix Λ_ψ of ψ is the following. Let q_i be a minimal idempotent in A_i and let z_j be the identity of B_j (a

minimal central idempotent in B). Then $\psi(q_i)z_j$ is the sum of $\Lambda_\psi(j, i)$ minimal idempotents in B_j .

Sometimes it is convenient to encode an inclusion matrix (or multiplicity matrix) by a bipartite graph, called the *branching diagram*; the branching diagram has vertices labeled by I arranged on one horizontal line, vertices labelled by J arranged along a second (higher) horizontal line, and $\Lambda_\psi(j, i)$ edges connecting $j \in J$ to $i \in I$.

If $A^{(1)} \subseteq A^{(2)} \subseteq A^{(3)} \dots$ is a sequence of inclusions of finite dimensional split semisimple algebras over F , then the branching diagram for the sequence is obtained by stacking the branching diagrams for each inclusion, with the upper vertices of the diagram for $A^{(i)} \subseteq A^{(i+1)}$ being identified with the lower vertices of the diagram for $A^{(i+1)} \subseteq A^{(i+2)}$.

Let $A \subseteq B$ be a pair of finite dimensional split semisimple algebras over F . The *Jones basic construction* for $A \subseteq B$ is the pair $B \subseteq \text{End}(B_A)$. That is, B is regarded as acting on itself by left multiplication, and is thus included in the endomorphisms of B as a right A -module. Then $\text{End}(B_A)$ is again a split semisimple algebra whose minimal ideals are in one-to-one correspondence with those of A , and the inclusion matrix for the pair $B \subseteq \text{End}(B_A)$ is the transpose of that for $A \subseteq B$.

Suppose now that B has a faithful trace ε with faithful restriction to A . Then there is a unique trace preserving conditional expectation (i.e. A - A bimodule map) $\varepsilon_A : B \rightarrow A$ determined by $\varepsilon(ba) = \varepsilon(\varepsilon_A(b)a)$ for all $a \in A$ and $b \in B$. Note that ε_A is an idempotent in $\text{End}(B_A)$. The Jones basic construction has a particular realization in terms of this data. In fact, $\text{End}(B_A) = \langle B, \varepsilon_A \rangle = B\varepsilon_AB$, where $\langle B, \varepsilon_A \rangle$ denotes the subalgebra of $\text{End}(B_A)$ generated by B and ε_A . Moreover, if q_i is a minimal idempotent in A_i , then $q_i\varepsilon_A = \varepsilon_A q_i$ is a minimal idempotent in the corresponding minimal ideal of $\text{End}(B_A)$.

Theorem 2.17 (Wenzl [29]). *Suppose $A \subseteq B \subseteq C$ are F -algebras with A and B finite dimensional split semisimple. Suppose that B has a trace $\varepsilon \in B^*$ that is non-degenerate on B and has non-degenerate restriction to A . Let ε_A denote the corresponding trace preserving conditional expectation $\varepsilon_A : B \rightarrow A$.*

Suppose that $e \in C$ is an idempotent such that $exe = \varepsilon_A(x)e$ for all $x \in B$ and $a \mapsto ae$ is an injective homomorphism from A to C . Let $\langle B, e \rangle$ be the subalgebra

of C generated by B and e . Then BeB is an ideal in $\langle B, e \rangle$, and

$$BeB \cong B\varepsilon_A B = \langle B, \varepsilon_A \rangle = \text{End}(B_A).$$

Remark 2.18. In the situation of the theorem, it follows that BeB is split semisimple, with minimal ideals in one-to-one correspondence with those of A . If q_i is a minimal idempotent in a minimal ideal A_i of A , then $q_i e$ is a minimal idempotent in the corresponding minimal ideal of BeB . BeB has an identity element z , necessarily a central idempotent in $\langle B, e \rangle$. The multiplicity matrix for the homomorphism $b \mapsto zb$ from B to BeB is the transpose of the inclusion matrix for $A \subseteq B$.

We now discuss *path idempotents* for a sequence $A^{(0)} \subseteq A^{(1)} \subseteq A^{(2)} \dots$ of split semisimple algebras, under the simplifying assumption (sufficient for our purposes) that $A^{(0)} = F$ and each inclusion $A^{(i)} \subseteq A^{(i+1)}$ is *multiplicity-free*, that is, the branching diagram has at most one edge connecting any pair of vertices. Let Γ_k be an index set for the minimal ideals of $A^{(k)}$; denote the minimal ideals of $A^{(k)}$ by $A_\lambda^{(k)}$ and the minimal central idempotents by $z_\lambda^{(k)}$ ($\lambda \in \Gamma_k$). Denote the unique element of Γ_0 by $\emptyset$. For $\mu \in \Gamma_j$ and $\lambda \in \Gamma_{j+1}$, write $\mu \subseteq \lambda$ if μ and λ are connected by an edge of the branching diagram. A *path* of length k in the branching diagram is a sequence $(\emptyset, \lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(k)})$ with $\lambda^{(j)} \subseteq \lambda^{(j+1)}$ for all j . Let $\mathcal{T}(k)$ be the set of all paths of length k , and $\mathcal{T}(k, \lambda)$ the set of paths of length k with final vertex λ . If $T = (\emptyset, \lambda^{(1)}, \dots, \lambda^{(k)})$ is a path of length k , let T' denote its truncation of length $k-1$, $T' = (\emptyset, \lambda^{(1)}, \dots, \lambda^{(k-1)})$. Moreover, if $\lambda \in \Gamma_{k+1}$ and $\lambda^{(k)} \subseteq \lambda$, let $T + \lambda$ denote the extension of T of length $k+1$ with final vertex λ . We claim that for each k , there is a canonical family of idempotents $\{p_T : T \in \mathcal{T}(k)\}$, in $A^{(k)}$, with the properties:

- (1) The elements p_T are mutually orthogonal minimal idempotents in $A^{(k)}$, and the sum of the p_T with $T \in \mathcal{T}(k, \lambda)$ is the minimal central idempotent $z_\lambda^{(k)}$.
- (2) $p_T p_{T'} = p_{T'} p_T = p_T$.

The construction is simple: Let $\emptyset$ denote the path of length 0. Necessarily $p_\emptyset = 1$. If the idempotents p_T have been constructed for paths of length j , $T \in \mathcal{T}(j, \mu)$, and $\lambda \in \Gamma_{j+1}$ satisfies $\mu \subseteq \lambda$, then put $p_{T+\lambda} = p_T z_\lambda^{(j+1)}$, which is a minimal idempotent in $A^{(j+1)}$ because of the assumption of multiplicity-free inclusions.

It follows that if $T = (\emptyset, \lambda^{(1)}, \lambda^{(2)}, \dots, \lambda^{(k)})$ is a path of length k , then for all $j \leq k$, $A^{(j)} p_T = A_{\lambda^{(j)}}^{(j)} p_T$ affords the simple $A_{\lambda^{(j)}}^{(j)}$ -module.

Lemma 2.19. *Suppose $A^{(0)} \subseteq A^{(1)} \subseteq A^{(2)} \subseteq \cdots \subseteq A^{(n)}$ is a sequence of split semisimple algebras, with $A^{(0)} = F$ and with each inclusion $A^{(i)} \subseteq A^{(i+1)}$ multiplicity-free. Suppose we have a trace ε on A_n , not necessarily faithful, but with faithful restriction to A_{n-1} . Suppose also that we have a trace-preserving conditional expectation $\varepsilon_n : A_n \rightarrow A_{n-1}$. Let T be a path of length n on the branching diagram for the sequence of inclusions. Then*

$$\varepsilon_n(p_T) = (\varepsilon(p_T)/\varepsilon(p_{T'}))p_{T'}.$$

Proof. We have $\varepsilon_n(p_T) = \varepsilon_n(p_{T'}p_Tp_{T'}) = p_{T'}\varepsilon_n(p_T)p_{T'}$. Since $p_{T'}$ is a minimal idempotent in A_{n-1} , we have $\varepsilon_n(p_T) = \kappa p_{T'}$ for some $\kappa \in F$. Taking traces, we have $\varepsilon(p_T) = \kappa\varepsilon(p_{T'})$, so $\kappa = \varepsilon(p_T)/\varepsilon(p_{T'})$. $\square$

3. REPRESENTATIONS OF THE 2-STRAND ALGEBRA AND ADMISSIBILITY CONDITIONS

To obtain any substantial results about the cyclotomic BMW and Kauffman tangle algebras, it appears to be necessary to impose a stronger condition on the ground ring than weak admissibility. Appropriate conditions can be found by considering the representation theory of the 2-strand algebra, in particular representations in which the generator e_1 acts non-trivially.

This idea was used by Ariki, Mathas, and Rui in a related context [3]; the analogue of their condition, translated to our context, is useful, but too restrictive. The “correct” notion of admissibility for cyclotomic BMW algebras was discovered by Wilcox and Yu [31]. The left module $W_{2,S,r}e_1$ is spanned over the ground ring by $\{e_1, y_1e_1, \dots, y_1^{r-1}e_1\}$, as is easily checked. Generically, one would expect this spanning set to be linearly independent over S , and the requirement of linear independence is the module-theoretic formulation of the admissibility condition of Wilcox and Yu.

3.1. Admissibility. Consider a ground ring S with parameters ρ, q, δ_j ($j \geq 0$) and $u_1, \dots, u_r$. Let a_j denote the signed elementary symmetric function in $u_1, \dots, u_r$, $a_j = (-1)^{r-j}\varepsilon_{r-j}(u_1, \dots, u_r)$. Let W_2 denote the cyclotomic BMW algebra $W_2 = W_{2,S,r}(u_1, \dots, u_r)$. Write e for e_1 and g for g_1 .

Lemma 3.1. *The left ideal W_2e in W_2 is equal to the S -span of $\{e, y_1e, \dots, y_1^{r-1}e\}$.*

Theorem 3.2 (Wilcox-Yu, [31]). *Let S be a ground ring with parameters ρ, q, δ_j ($j \geq 0$) and $u_1, \dots, u_r$. Assume that $(q - q^{-1})$ is non-zero and not a zero-divisor in S . The following conditions are equivalent:*

- (1) *S is weakly admissible, and $\{e, y_1 e, \dots, y_1^{r-1} e\} \subseteq W_2$ is linearly independent over S .*
- (2) *The parameters satisfy the following relations:*

$$(3.1) \quad \rho(a_\ell - a_{r-\ell}/a_0) + (q - q^{-1}) \left[\sum_{j=1}^{r-\ell} a_{j+\ell} \delta_j - \sum_{j=\max(\ell+1, \lceil r/2 \rceil)}^{\lfloor (\ell+r)/2 \rfloor} a_{2j-\ell} + \sum_{j=\lceil \ell/2 \rceil}^{\min(\ell, \lceil r/2 \rceil - 1)} a_{2j-\ell} \right] = 0, \\ \text{for } 1 \leq \ell \leq r-1,$$

$$(3.2) \quad \rho^{-1} a_0 - \rho a_0^{-1} = \begin{cases} 0 & \text{if } r \text{ is odd} \\ (q - q^{-1}) & \text{if } r \text{ is even,} \end{cases}$$

and

$$(3.3) \quad \delta_a = - \sum_{j=0}^{r-1} a_j \delta_{a-r+j} \quad \text{for } a \geq r.$$

- (3) *S is weakly admissible, and W_2 admits a module M with an S -basis $\{v_0, y_1 v_0, \dots, y_1^{r-1} v_0\}$ such that $ev_0 = \delta_0 v_0$.*

Definition 3.3 (Wilcox and Yu, [31]). Let S be a ground ring with parameters ρ, q, δ_j ($j \geq 0$) and $u_1, \dots, u_r$. One says that S is *admissible* (or that the parameters are *admissible*) if $(q - q^{-1})$ is non-zero and not a zero-divisor in S and if the equivalent conditions of Theorem 3.2 hold.

In the following statement, we omit the assumption that $q - q^{-1}$ is not a zero-divisor, but we impose the assumption that S is weakly admissible. The statement can be proved by a minor modification of the proof of Wilcox and Yu.

Corollary 3.4. *Let S be a weakly admissible ring with parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. The following conditions are equivalent.*

- (1) *$\{e, y_1 e, \dots, y_1^{r-1} e\} \subseteq W_2$ is linearly independent over S .*
- (2) *The parameters satisfy the relations (3.1) and (3.2) of Wilcox and Yu.*
- (3) *W_2 admits a cyclic module M with an S -basis $\{v_0, y_1 v_0, \dots, y_1^{r-1} v_0\}$ such that $ev_0 = \delta_0 v_0$.*

Remark 3.5.

- (1) Let S be a ground ring with admissible parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. Then

$$\rho, -q^{-1}, \delta_j \ (j \geq 0), \text{ and } u_1, \dots, u_r$$

and

$$-\rho, -q, \delta_j \ (j \geq 0), \text{ and } u_1, \dots, u_r$$

are also sets of admissible parameters. This follows from condition (2) of Theorem 3.2.

- (2) If S is a ground ring with admissible parameters and $\varphi : S \rightarrow S'$ is a morphism of ground rings in the sense of Definition 2.13, such that $\varphi(q - q^{-1})$ is non-zero and not a zero divisor, then S' is also admissible.

Remark 3.6. Assume that S is a ground ring with admissible parameters. We observe that the admissibility relation (3.1) can be solved for

$$(q - q^{-1})\delta_1, \dots, (q - q^{-1})\delta_{r-1}$$

in terms of $\rho, a_0^{\pm 1}, a_1, \dots, a_{r-1}$, and $q^{\pm 1}$. Rewrite the relations (3.1) in the form

$$(3.4) \quad (q - q^{-1}) \sum_{j=1}^{r-\ell} a_{j+\ell} \delta_j = p_\ell(q^{\pm 1}, \rho, a_0^{\pm 1}, a_1, \dots, a_{r-1}) \quad (1 \leq \ell \leq r-1),$$

where the p_ℓ are specific polynomials. This is a triangular system of linear equations in the variables $(q - q^{-1})\delta_j$ with a unitriangular matrix of coefficients

$$\begin{bmatrix} 1 & & & & \\ a_{r-1} & 1 & & & \\ a_{r-2} & a_{r-1} & 1 & & \\ \vdots & & \ddots & \ddots & \\ a_2 & a_3 & \dots & a_{r-1} & 1 \end{bmatrix},$$

so it has a solution of the form

$$(q - q^{-1})\delta_j = Q_j(q^{\pm 1}, \rho, a_0^{\pm 1}, a_1, \dots, a_{r-1}), \quad \text{for } 1 \leq j \leq r-1,$$

where the Q_j are again some specific polynomials. Because $q - q^{-1}$ is assumed not to be a zero-divisor in S , we can append an inverse of $q - q^{-1}$ to S . In $S[(q - q^{-1})^{-1}]$, we have

$$\delta_j = (q - q^{-1})^{-1} Q_j(q^{\pm 1}, \rho, a_0^{\pm 1}, a_1, \dots, a_{r-1}), \quad \text{for } 1 \leq j \leq r-1,$$

Remark 3.7. Equation (3.2) is equivalent to

$$\begin{cases} (a_0 - \rho)(a_0 + \rho) = 0 & \text{if } r \text{ is odd, and} \\ (\rho - q^{-1}a_0)(\rho + qa_0) = 0 & \text{if } r \text{ is even.} \end{cases}$$

If S is an integral domain, we have $\rho = \pm a_0$ if r is odd, and $\rho = -qa_0$ or $\rho = q^{-1}a_0$ if r is even. However, we do not need to entertain both solutions. Considering Remark 3.5, we can assume

$$\rho = -a_0 = \prod_{j=1}^r u_j,$$

if r is odd, and

$$\rho = q^{-1}a_0 = q^{-1} \prod_{j=1}^r u_j,$$

if r is even.

We have the following consequence of this discussion:

Corollary 3.8. *Let S be an integral ground ring with admissible parameters ρ , q , etc. Then $\rho^{\pm 1}$ and δ_j for $j \geq 0$ are elements of the subring of $S[(q - q^{-1})^{-1}]$ generated by $q^{\pm 1}$, $(q^{-1} - q)^{-1}$, and $u_1^{\pm 1}, \dots, u_r^{\pm 1}$.*

3.2. u -admissibility. We maintain the notation established at the beginning of Section 3.1. In this section, we assume that the ring S is an integral domain, that $q - q^{-1} \neq 0$, that the parameters u_i are distinct, and that $u_i u_j \neq 1$ for all i, j .

Lemma 3.9. *Let F be field and let $u_1, \dots, u_r$ be distinct non-zero elements of F with $1 \neq u_i u_j$ for all i, j . Let ρ and q be non-zero elements of F with $q^2 \neq 1$. Then the unique solution to the system of linear equations*

$$(3.5) \quad \sum_j \frac{1}{1 - u_i u_j} \gamma_j = \frac{1}{1 - u_i^2} + \frac{1}{\rho(q^{-1} - q)} \quad (1 \leq i \leq r)$$

is

$$(3.6) \quad \gamma_j = \prod_{\ell \neq j} \frac{(u_\ell u_j - 1)}{u_j - u_\ell} \left(\frac{1 - u_j^2}{\rho(q^{-1} - q)} \prod_{\ell \neq j} u_\ell + \begin{cases} 1 & \text{if } r \text{ is odd} \\ -u_j & \text{if } r \text{ is even} \end{cases} \right)$$

We defer the proof of the lemma to Remark 3.16 in Section 3.3.

Theorem 3.10. *Let S be an integral ground ring with parameters ρ , q , δ_j ($j \geq 0$) and $u_1, \dots, u_r$. Assume that $(q - q^{-1}) \neq 0$, that the elements u_i are distinct, and that $u_i u_j \neq 1$ for all i, j . Define γ_j in the field of fractions of S by (3.6), for $1 \leq j \leq r$.*

The following conditions are equivalent:

- (1) S is weakly admissible, and $\{e, y_1 e, \dots, y_1^{r-1} e\} \subseteq W_{2,S}$ is linearly independent over S .
- (2) For all $a \geq 0$, we have $\delta_a = \sum_{j=1}^r \gamma_j u_j^a$.
- (3) S is weakly admissible, and $W_{2,S}$ admits a module M with an S -basis $\{v_0, y_1 v_0, \dots, y_1^{r-1} v_0\}$ such that $ev_0 = \delta_0 v_0$.

Proof. Let F denote the field of fractions of S , and let (1_F) and (3_F) indicate the analogue of conditions (1) and (3) over F , i.e.

(1_F) F is weakly admissible, and $\{e, y_1 e, \dots, y_1^{r-1} e\} \subseteq W_{2,F}$ is linearly independent over F .

(3_F) F is weakly admissible, and $W_{2,F}$ admits a module M with an F -basis $\{v_0, y_1 v_0, \dots, y_1^{r-1} v_0\}$ such that $ev_0 = \delta_0 v_0$.

According to Theorem 3.2, conditions (1) and (3) are equivalent, and moreover, they are equivalent to the second condition of Theorem 3.2. Since that condition is unchanged if we replace S by F , it follows that (1) and (3) are equivalent to (1_F) and (3_F) . Therefore, to prove the theorem, it suffices to show that

$$(1_F) \implies (2) \implies (3_F).$$

Assume condition (1_F) . Recall that the span of $\{e, y_1 e, \dots, y_1^{r-1} e\}$ is equal to the left ideal $W_{2,F} e$. Moreover, $ey_1^j e = \delta_j e$, so $eW_{2,F} e = Fe$.

For each j ($1 \leq j \leq r$) define p_j by

$$p_j = \prod_{i \neq j} \frac{y_1 - u_i}{u_j - u_i} \in W_{1,F}.$$

Then we have $y_1 p_j = u_j p_j$, $p_j^2 = p_j$, and $\sum_j p_j = 1$.

Let $m_j = p_j e$. Then $m_j \neq 0$, by the linear independence of $\{e, y_1 e, \dots, y_1^{r-1} e\}$ over F , $y_1 m_j = u_j m_j$, and $\sum_j m_j = e$. The set $\{m_1, \dots, m_r\}$ is linearly independent, since the m_j are eigenvectors for y_1 with distinct eigenvalues. Because the dimension of $W_{2,F} e$ is r , $\{m_1, \dots, m_r\}$ is a basis of $W_{2,F} e$.

Now we define $\kappa_j \in F$ by $em_j = \kappa_j e = \kappa_j \sum_i m_i$ and $c_{i,j}$ by $gm_j = \sum_i c_{i,j} m_i$. Note that $y_1 g(y_1^s e) = g^{-1} y_1^{-1} (y_1^s e)$, by Remark 2.4. Therefore,

$$y_1 g m_j = g^{-1} y_1^{-1} m_j = \left(g + (q - q^{-1})(e - 1) \right) y_1^{-1} m_j.$$

We have

$$y_1 g m_j = y_1 \sum_i c_{i,j} m_i = \sum_i u_i c_{i,j} m_i.$$

On the other hand,

$$\begin{aligned} & (g + (q - q^{-1})(e - 1))y_1^{-1}m_j \\ &= u_j^{-1} \sum_i c_{i,j} m_i + u_j^{-1}(q - q^{-1})(\kappa_j \sum_i m_i - m_j) \end{aligned}$$

Equating the coefficient of m_i in the two expressions gives

$$(u_i - u_j^{-1})c_{i,j} = (q - q^{-1})u_j^{-1}(\kappa_j - \delta(i, j)),$$

where $\delta(i, j)$ denotes the Kronecker delta. Solving, we get

$$(3.7) \quad c_{i,j} = (q^{-1} - q) \frac{\kappa_j - \delta(i, j)}{1 - u_i u_j}.$$

If we apply both sides of the equation $ge = \rho^{-1}e$ to m_k , we get

$$\kappa_k \sum_{i,j} c_{i,j} m_i = \rho^{-1} \kappa_k \sum_i m_i.$$

Choosing k so that $\kappa_k \neq 0$, and equating the coefficients of m_i on both sides gives $\sum_j c_{i,j} = \rho^{-1}$ for all i . Taking (3.7) into account, we have for all i ,

$$(3.8) \quad \sum_j \frac{1}{1 - u_i u_j} \kappa_j = \frac{1}{1 - u_i^2} + \frac{1}{\rho(q^{-1} - q)}.$$

It follows from (3.8) and Corollary 3.9 that κ_j is given by

$$(3.9) \quad \kappa_j = \gamma_j = \prod_{\ell \neq j} \frac{(u_\ell u_j - 1)}{u_j - u_\ell} \left(\frac{1 - u_j^2}{\rho(q^{-1} - q)} \prod_{\ell \neq j} u_\ell + \begin{cases} 1 & \text{if } r \text{ is odd} \\ -u_j & \text{if } r \text{ is even} \end{cases} \right)$$

Next, note that for $a \geq 0$

$$\delta_a e = e y_1^a e = e y_1^a \sum_i m_i = e \sum_i u_i^a m_i = \left(\sum_i \gamma_i u_i^a \right) e.$$

Therefore, for $a \geq 0$, $\delta_a = \sum_i \gamma_i u_i^a$. This completes the proof of the implication $(1_F) \implies (2)$.

We now assume condition (2) and prove (3_F). Let $\{\hat{e}_1, \dots, \hat{e}_r\}$ denote the standard basis of $M = F^r$. Define linear maps E, G, Y on M by

$$(3.10) \quad \begin{aligned} Y \hat{e}_j &= u_j \hat{e}_j, \\ E \hat{e}_j &= \gamma_j (\hat{e}_1 + \dots + \hat{e}_r) \\ G \hat{e}_j &= (q^{-1} - q) \sum_{i=1}^r \frac{\gamma_j - \delta(i, j)}{1 - u_i u_j} \hat{e}_i, \end{aligned}$$

where for $1 \leq j \leq r$, γ_j is given by (3.6).

Applying the definitions of Y, G, E , and the u -admissibility assumption, we have, for $a \geq 0$, $EY^a E \hat{e}_j = (\sum_i \gamma_i u_i^a) E \hat{e}_j = \delta_a E \hat{e}_j$. Thus, for $a \geq 0$,

$$(3.11) \quad EY^a E = \delta_a E,$$

and, in particular, $E^2 = \delta_0 E$. It is straightforward to compute that

$$(3.12) \quad \begin{aligned} YGY &= G + (q^{-1} - q)(1 - E), \text{ and} \\ GE &= EG = \rho^{-1} E. \end{aligned}$$

We give some details for the computation of YGY ; applying YGY to $\hat{e}_j$ gives

$$YGY \hat{e}_j = (q^{-1} - q) \sum_k \frac{u_j u_k (\gamma_j - \delta(j, k))}{1 - u_j u_k} \hat{e}_k.$$

If we write $u_j u_k = (u_j u_k - 1) + 1$ in the numerator and simplify, we get

$$\begin{aligned} & (q^{-1} - q)(-\gamma_j \sum_k \hat{e}_k + \hat{e}_j) + (q^{-1} - q) \sum_k \frac{(\gamma_j - \delta(j, k))}{1 - u_j u_k} \hat{e}_k \\ &= (q^{-1} - q)(-E + 1) \hat{e}_j + G \hat{e}_j. \end{aligned}$$

Taking into account (3.12), we see that YGY commutes with both E and G ,

$$(3.13) \quad \begin{aligned} GYGY &= YGYG, \text{ and} \\ EYGY &= YGYE. \end{aligned}$$

Similarly, we compute

$$YGYE \hat{e}_j = (q^{-1} - q) \gamma_j \sum_k \sum_i \frac{u_i u_k (\gamma_i - \delta(i, k))}{1 - u_i u_k} \hat{e}_k.$$

Simplifying this as above, we get

$$(q^{-1} - q) \gamma_j (-\sum_i \gamma_i + 1) \sum_k \hat{e}_k + (q^{-1} - q) \gamma_j \sum_k \sum_i \frac{(\gamma_i - \delta(i, k))}{1 - u_i u_k} \hat{e}_k.$$

Using $\sum_i \gamma_i = \delta_0$ and (3.5), we can simplify this to

$$((q^{-1} - q)(-\delta_0 + 1) + \rho^{-1}) \gamma_j \sum_k \hat{e}_k = \rho \gamma_j \sum_k \hat{e}_k = \rho E \hat{e}_j$$

Thus, we have

$$(3.14) \quad EYGY = YGYE = \rho E.$$

Set $m = \sum_i \hat{e}_i$. We claim that $\{m, Ym, \dots, Y^{r-1}m\}$ is a basis of M . In fact, if $\sum_{j=0}^{r-1} \alpha_j Y^j m = 0$, then $\sum_{i=1}^r (\sum_{j=0}^{r-1} \alpha_j u_i^j) \hat{e}_i = 0$. Since the u_i are distinct, the Vandermonde matrix (u_i^j) is invertible, so $\alpha_j = 0$ for all j . This shows that $\{m, Ym, \dots, Y^{r-1}m\}$ is linearly independent, and therefore a basis of M .

We claim that $GYGY$ is the identity transformation. By the last paragraph, it is enough to show that $GYGY(Y^k m) = Y^k m$ for $0 \leq k \leq r-1$. Note that $YGYGE = E$, by (3.12) and (3.14), and also $Em = \delta_0 m$. According to (3.13), Y commutes with GYG , so

$$\begin{aligned} GYGY(Y^k m) &= Y^k YGYGm = (1/\delta_0) Y^k YGYGE m \\ &= (1/\delta_0) Y^k Em = Y^k m. \end{aligned}$$

Using (3.12), we have

$$(3.15) \quad G^{-1} = YGY = G + (q^{-1} - q)(1 - E).$$

It follows from (3.11)–(3.15) that E, G, Y satisfy the relations of the cyclotomic BMW algebra with parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. In particular, since we are working over a field, and $E \neq 0$, the parameters must be weakly admissible; see the statement just before Definition 2.14. $\square$

Definition 3.11. Let S be an integral ground ring with parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. Say that the parameters are *u-admissible* or that S is *u-admissible* if:

- (1) $q - q^{-1} \neq 0$, $u_i \neq u_j$ for $i \neq j$ and $u_i u_j \neq 1$ for any i, j , and
- (2) the equivalent conditions of Theorem 3.10 hold.

Corollary 3.12. Let S be an integral ground ring with parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. Then S is *u-admissible* if, and only if, S is admissible, the u_i are distinct and $u_i u_j \neq 1$ for all i, j .

Proof. This follows from Theorems 3.2 and 3.10. $\square$

Remark 3.13. Let S be a ground ring with *u-admissible* parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. Then

$$\rho, -q^{-1}, \delta_j \ (j \geq 0), \text{ and } u_1, \dots, u_r$$

and

$$-\rho, -q, \delta_j \ (j \geq 0), \text{ and } u_1, \dots, u_r$$

are also sets of *u-admissible* parameters. Moreover, the quantities γ_j computed from any of these sets of parameters according to (3.6) are the same.

Remark 3.14. Let S be an integral ground ring with *u-admissible* parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. By Remarks 3.7 and 3.13, we can assume $\rho = \prod_j u_j$

if r is odd and $\rho = q^{-1} \prod_j u_j$ if r is even. Using this, we can rewrite the formula for γ_j (3.6) as

$$(3.16) \quad \gamma_j = \frac{\rho(u_j - q^{-1})(u_j + q)}{u_j^2(q - q^{-1})} \prod_{\ell \neq j} \frac{u_j - u_\ell^{-1}}{u_j - u_\ell} \quad \text{if } r \text{ is odd,}$$

and

$$(3.17) \quad \gamma_j = \frac{\rho(u_j - q)(u_j + q)}{u_j^2(q - q^{-1})} \prod_{\ell \neq j} \frac{u_j - u_\ell^{-1}}{u_j - u_\ell} \quad \text{if } r \text{ is even.}$$

Note that if $u_j \notin \{\pm q^{\pm 1}\}$, then $\gamma_j \neq 0$.

3.3. Some generating functions. We will investigate some generating functions that appear naturally in the study of cyclotomic BMW algebras. Let $u_1, \dots, u_r, q$, and ρ be indeterminants over $\mathbb{Z}$. We will need to consider rational functions γ_j in these variables that are solutions to the system of linear equations:

$$\sum_j \frac{1}{1 - u_i u_j} \gamma_j = \frac{1}{1 - u_i^2} + \frac{1}{\rho(q^{-1} - q)} \quad (1 \leq i \leq r).$$

Lemma 3.15. *Let F be field and let $u_1, \dots, u_r$ be distinct non-zero elements of F with $1 \neq u_i u_j$ for all i, j .*

(1) *The matrix*

$$\left(\frac{1}{1 - u_i u_j} \right)_{1 \leq i, j \leq r}$$

is invertible.

(2) *The unique solution to the linear system of equations*

$$\sum_j \frac{1}{1 - u_i u_j} \gamma_j^{(1)} = 1 \quad (1 \leq i \leq r)$$

is

$$\gamma_j^{(1)} = (1 - u_j^2) \prod_{\ell \neq j} u_\ell \prod_{\ell \neq j} \frac{(u_\ell u_j - 1)}{u_j - u_\ell}.$$

(3) *The unique solution to the linear system of equations*

$$\sum_j \frac{1}{1 - u_i u_j} \gamma_j^{(2)} = \frac{1}{1 - u_i^2} \quad (1 \leq i \leq r)$$

is

$$\gamma_j^{(2)} = \prod_{\ell \neq j} \frac{u_\ell u_j - 1}{u_j - u_\ell} \quad \text{if } r \text{ is odd,}$$

and

$$\gamma_j^{(2)} = -u_j \prod_{\ell \neq j} \frac{u_\ell u_j - 1}{u_j - u_\ell} \quad \text{if } r \text{ is even.}$$

Proof. Recall Cauchy's determinant identity [15], Section 2.1,

$$\det \left(\frac{1}{1 - x_i y_j} \right) = \frac{\prod_{i < j} (x_i - x_j)(y_i - y_j)}{\prod_{i,j} (1 - x_i y_j)},$$

which specializes to

$$\det \left(\frac{1}{1 - u_i u_j} \right) = \frac{\prod_{i < j} (u_i - u_j)^2}{\prod_{i,j} (1 - u_i u_j)}.$$

Since $u_i \neq u_j$ for $i \neq j$, this implies statement (1).

For part (2), we have to prove the identity, for all i ($1 \leq i \leq r$):

$$\sum_j \frac{1}{1 - u_i u_j} (1 - u_j^2) \prod_{\ell \neq j} u_\ell \prod_{\ell \neq j} \frac{(u_\ell u_j - 1)}{u_j - u_\ell} = 1,$$

or

$$(3.18) \quad \sum_j \prod_{\ell \neq i} (u_\ell u_j - 1) \prod_{\ell \neq j} \frac{u_\ell}{u_j - u_\ell} = 1.$$

By the principal of permanence of identities (see [4], p. 456) it suffices to prove this when the u_i are distinct non-zero complex numbers satisfying $u_i u_j \neq 1$ for all i, j . Put

$$f_i(\zeta) = (1/\zeta) \prod_{\ell \neq i} (\zeta u_\ell - 1) \prod_{\ell=1}^r \frac{u_\ell}{\zeta - u_\ell}.$$

Check that the residue of $f_i(\zeta)$ at $\zeta = u_j$ is the summand on the left side of (3.18). The other possible poles of $f_i(\zeta)$ are at $\zeta = 0$ and at $\zeta = \infty$. The residue at $\zeta = 0$ is -1 and the residue at $\zeta = \infty$ is zero. The identity of (3.18) follows, since the sum of the residues of $f_i(\zeta)$ at its finite poles and at ∞ is zero.

For part (3), we have to prove the identity for all i ($1 \leq i \leq r$):

$$(3.19) \quad \sum_j \frac{1 - u_i^2}{1 - u_i u_j} \prod_{\ell \neq j} \frac{u_\ell u_j - 1}{u_j - u_\ell} = 1 \quad \text{if } r \text{ is odd,}$$

and

$$(3.20) \quad \sum_j \frac{1 - u_i^2}{1 - u_i u_j} (-u_j) \prod_{\ell \neq j} \frac{u_\ell u_j - 1}{u_j - u_\ell} = 1 \quad \text{if } r \text{ is even.}$$

Again, we may assume that the u_i are distinct non-zero complex numbers. When r is odd, we put

$$f_i(\zeta) = \frac{1 - u_i^2}{(\zeta^2 - 1)(1 - \zeta u_i)} \prod_{\ell=1}^r \frac{\zeta u_\ell - 1}{\zeta - u_\ell},$$

and observe that the residue of $f_i(\zeta)$ at $\zeta = u_j$ is the summand on the left side of (3.19). The other possible poles of $f_i(\zeta)$ are at $\zeta = \pm 1$ and $\zeta = \infty$. The residue at $\zeta = 1$ is $(-1/2)(1 + u_i)$, the residue at $\zeta = -1$ is $(-1/2)(1 - u_i)$, and the residue at $\zeta = \infty$ is zero. Since the sum of these residues is -1 , Equation (3.19) follows. The proof of (3.20) is similar, and we omit it. $\square$

Remark 3.16. Lemma 3.9 from Section 3.2 is a corollary of Lemma 3.15. Namely, the unique solution to the system of equations

$$\sum_j \frac{1}{1 - u_i u_j} \gamma_j = \frac{1}{1 - u_i^2} + \frac{1}{\rho(q^{-1} - q)} \quad (1 \leq i \leq r)$$

is

$$\gamma_j = \frac{1}{\rho(q^{-1} - q)} \gamma_j^{(1)} + \gamma_j^{(2)}.$$

Fix a natural number r . Let $u_1, \dots, u_r$ and t be indeterminants over $\mathbb{Z}$, and define

$$(3.21) \quad G(t) = G(u_1, \dots, u_r; t) = \prod_{\ell=1}^r \frac{t - u_\ell}{t u_\ell - 1}.$$

Let $\mu_a = \mu_a(u_1, \dots, u_r)$ denote the coefficients of the formal power series expansion of $G(t)$,

$$(3.22) \quad G(t) = \sum_{a=0}^{\infty} \mu_a t^a.$$

Notice that each μ_a is a symmetric polynomial in $u_1, \dots, u_r$ and that $G(t^{-1}) = G(t)^{-1}$.

Now suppose that ρ and q are additional indeterminants, and for $1 \leq j \leq r$, define rational functions γ_j by (3.6). Moreover, for $a \geq 0$, define $\delta_a = \sum_{j=1}^r \gamma_j u_j^a$. Let $Z_1(t)$ be the generating function for the δ_a ,

$$(3.23) \quad \begin{aligned} Z_1(t) &= \sum_{a \geq 0} \delta_a t^{-a} = \sum_{a \geq 0} \sum_{j=1}^r \gamma_j u_j^a t^{-a} \\ &= \sum_{j=1}^r \gamma_j \sum_{a \geq 0} u_j^a t^{-a} = \sum_{j=1}^r \gamma_j \frac{t}{t - u_j}. \end{aligned}$$

In the following, we use the notation

$$\delta(P) = \begin{cases} 1 & \text{if } P \text{ is true} \\ 0 & \text{if } P \text{ is false.} \end{cases}$$

Lemma 3.17. *Let p denote $\prod_{j=1}^r u_j$.*

$$(1) \quad Z_1(t) = \frac{1}{\rho(q^{-1} - q)} + \frac{t^2}{t^2 - 1} + A(t) G(t^{-1}), \text{ where}$$

$$A(t) = \begin{cases} \rho^{-1}p/(q - q^{-1}) + t/(t^2 - 1) & \text{if } r \text{ is odd, and} \\ \rho^{-1}p/(q - q^{-1}) - t^2/(t^2 - 1) & \text{if } r \text{ is even.} \end{cases}$$

(2) *If r is odd, then for $a \geq 0$,*

$$\begin{aligned} \delta_a = & \delta(a = 0) \rho^{-1}/(q - q^{-1}) + \delta(a \text{ is even}) \\ & + \mu_a \rho^{-1}p/(q - q^{-1}) + \mu_{a-1} + \mu_{a-3} + \cdots \end{aligned}$$

(3) *If r is even, then for $a \geq 0$,*

$$\begin{aligned} \delta_a = & \delta(a = 0) \rho^{-1}/(q - q^{-1}) + \delta(a \text{ is even}) \\ & + \mu_a \rho^{-1}p/(q - q^{-1}) - \mu_a - \mu_{a-2} - \mu_{a-4} + \cdots \end{aligned}$$

$$(4) \quad \delta_0 = \frac{1 - p^2}{\rho(q^{-1} - q)} + 1 - \begin{cases} 0 & \text{if } r \text{ is odd} \\ p & \text{if } r \text{ is even.} \end{cases}$$

(5) *For all $a \geq 0$, δ_a is an element of the ring*

$$\mathbb{Z}[u_1, \dots, u_r, \rho^{-1}, (q^{-1} - q)^{-1}],$$

and is symmetric in $u_1, \dots, u_r$.

Proof. To prove the identity of part (1), it suffices to suppose that the quantities $u_1, \dots, u_r$, t , q , and ρ are complex numbers, algebraically independent over $\mathbb{Q}$. Set

$$h(\zeta) = G(\zeta^{-1}) \left(\zeta^{-1} \frac{p}{\rho(q - q^{-1})} + \begin{cases} 1/(\zeta^2 - 1) & \text{if } r \text{ is odd} \\ -\zeta/(\zeta^2 - 1) & \text{if } r \text{ is even} \end{cases} \right).$$

It is straightforward to check that the residue of $h(\zeta)t/(t - \zeta)$ at $\zeta = u_j$ is

$$\gamma_j t/(t - u_j).$$

Thus, $Z_1(t)$ is the sum of the residues of $h(\zeta)t/(t - \zeta)$ at $u_1, \dots, u_r$. The other possible poles of $h(\zeta)t/(t - \zeta)$ are at $0, \pm 1, \infty$, and t . Independent of the parity of r , the residue of $h(\zeta)t/(t - \zeta)$ at $\zeta = 0$ is $1/(\rho(q - q^{-1}))$, the residue at $\zeta = 1$ is $-(1/2)(t/(t - 1))$, the residue at $\zeta = -1$ is $-(1/2)(t/(t + 1))$, and the residue at ∞ is zero. For r odd, the residue of $h(\zeta)t/(t - \zeta)$ at $\zeta = t$ is

$$G(t^{-1}) \left(p/(\rho(q^{-1} - q)) - t/(t^2 - 1) \right).$$

For r even, the residue of $h(\zeta)t/(t - \zeta)$ at $\zeta = t$ is

$$G(t^{-1}) \left(p/(\rho(q^{-1} - q)) + t^2/(t^2 - 1) \right).$$

$Z_1(t)$ is the negative of the sum of the residues of $h(\zeta)t/(t - \zeta)$ at $0, \pm 1, \infty$, and t . This yields the formula of part (1).

For part (2), write

$$\begin{aligned} Z_1(t^{-1}) &= \sum_{a \geq 0} \delta_a t^a = \frac{1}{\rho(q^{-1} - q)} + \frac{1}{1 - t^2} \\ &\quad + G(t) \left(\frac{p}{\rho(q - q^{-1})} + \begin{cases} t/(1 - t^2) & \text{if } r \text{ is odd} \\ -1/(1 - t^2) & \text{if } r \text{ is even} \end{cases} \right). \end{aligned}$$

One can now read off the coefficient of t^a on the right hand side.

Part (3) follows from part (2), using that $\mu_0 = G(0) = p$, and part (4) follows from part (2) as well. $\square$

Note that if F is a field with u -admissible parameters ρ, q, δ_j , and $u_1, \dots, u_r$, and $Z_1(t) = \sum_{a \geq 0} \delta_a t^{-a}$, then Lemma 3.17 applies, but we also can assume that $\rho = p$ if r is odd, and $\rho = q^{-1}p$ if r is even. So the formulas of the lemma simplify as follows:

$$\begin{aligned} (3.24) \quad Z_1(t) &= \frac{1}{\rho(q^{-1} - q)} + \frac{t^2}{t^2 - 1} + A(t) G(t^{-1}), \quad \text{where} \\ A(t) &= \begin{cases} 1/(q - q^{-1}) + t/t^2 - 1 & \text{if } r \text{ is odd, and} \\ q/(q - q^{-1}) - t^2/t^2 - 1 & \text{if } r \text{ is even.} \end{cases} \end{aligned}$$

3.4. The universal admissible ring. We begin by constructing a universal u -admissible integral domain. Let $\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r$ be algebraically independent indeterminates over $\mathbb{Z}$. Let S_0 be the Laurent polynomial ring in $\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r$, with $(\mathbf{q}^{-1} - \mathbf{q})^{-1}$ adjoined,

$$S_0 = \mathbb{Z}[\mathbf{q}^{\pm 1}, \mathbf{u}_1^{\pm 1}, \dots, \mathbf{u}_r^{\pm 1}][(\mathbf{q}^{-1} - \mathbf{q})^{-1}].$$

Let $\mathbf{p} = \prod_j \mathbf{u}_j$. If r is odd, put $\boldsymbol{\rho} = \mathbf{p}$, and if r is even, put $\boldsymbol{\rho} = \mathbf{q}^{-1}\mathbf{p}$. Define γ_j for $1 \leq j \leq r$ (in the field of fractions of S_0) by (3.6), and set $\boldsymbol{\delta}_a = \sum_{j=1}^r \gamma_j \mathbf{u}_j^a$. According to Lemma 3.17, $\boldsymbol{\delta}_a \in S_0$, and, moreover,

$$\boldsymbol{\delta}_0 = 1 + \frac{1 - \mathbf{p}^2}{\boldsymbol{\rho}(\mathbf{q}^{-1} - \mathbf{q})} - \begin{cases} 0 & \text{if } r \text{ is odd} \\ \mathbf{p} & \text{if } r \text{ is even.} \end{cases}$$

Taking into account that $\boldsymbol{\rho} = \mathbf{p}$ or $\boldsymbol{\rho} = \mathbf{q}^{-1}\mathbf{p}$, we have $\boldsymbol{\delta}_0 \neq 0$ by the algebraic independence of $\mathbf{q}$ and $\mathbf{u}_1, \dots, \mathbf{u}_r$. Moreover, one can check that

$\rho^{-1} - \rho = (q^{-1} - q)(\delta_0 - 1)$. Finally, define $\overline{S} = S_0[\delta_0^{-1}]$. Then $\overline{S}$ is u -admissible. Note that the field of fractions of $\overline{S}$ is $\mathbb{Q}(\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r)$.

Let S be another u -admissible integral domain with parameters ρ, q , etc. Using Remark 3.13, we can assume that $\rho = \prod_j u_j$ if r is odd and $\rho = q^{-1} \prod_j u_j$ if r is even. Because of the algebraic independence of $\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r$, there is a ring homomorphism $\varphi : \overline{S} \rightarrow S[(q - q^{-1})^{-1}]$ mapping $\mathbf{q} \mapsto q$ and $\mathbf{u}_i \mapsto u_i$ for each i . It follows that $\varphi : \rho \mapsto \rho$. Since $\overline{S}$ and S are u -admissible, we have $\delta_a \mapsto \delta_a$ for $a \geq 0$ as well.

We have shown the following result:

Lemma 3.18. *There is a “universal” integral domain $\overline{S}$ with u -admissible parameters $\rho, \mathbf{q}, \delta_j$, ($j \geq 0$), $\mathbf{u}_1, \dots, \mathbf{u}_r$, with the properties:*

- (1) $\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r$ are algebraically independent over $\mathbb{Z}$.
- (2) $\overline{S}$ is generated as a unital ring by $\mathbf{q}^{\pm 1}, (q^{-1} - q)^{-1}, \mathbf{u}_1^{\pm 1}, \dots, \mathbf{u}_r^{\pm 1}$, and δ_0^{-1} .
- (3) The field of fractions of $\overline{S}$ is $\mathbb{Q}(\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r)$.
- (4) Whenever S is an integral domain with u -admissible parameters ρ, q , etc., there exists a morphism of ground rings (see Definition 2.13)

$$\phi : \overline{S} \rightarrow S[(q - q^{-1})^{-1}].$$

Next we consider the construction of universal admissible rings. Let $\mathbf{q}, \rho, \delta_0, \dots, \delta_{r-1}, \mathbf{u}_1, \dots, \mathbf{u}_r$ be indeterminants over $\mathbb{Z}$. Let $\mathbf{a}_j$ be the signed elementary symmetric polynomials in the variables $\mathbf{u}_i$. For $a \geq r$, define δ_a by the recursion $\delta_a = -\sum_{j=0}^{r-1} \mathbf{a}_j \delta_{a-r+j}$.

Set

$$\mathbb{A} = \mathbb{Z}[\mathbf{q}^{\pm 1}, \rho^{\pm 1}, \delta_0^{\pm 1}, \delta_1, \dots, \delta_{r-1}, \mathbf{u}_1^{\pm 1}, \dots, \mathbf{u}_r^{\pm 1}].$$

If r is odd, set $f_+ = \rho + \mathbf{a}_0$ and $f_- = \rho - \mathbf{a}_0$. If r is even, set $f_+ = \rho - q^{-1} \mathbf{a}_0$ and $f_- = \rho + q \mathbf{a}_0$. For r of either parity, let $f_0 = f_+ f_-$. For $\varepsilon \in \{0, +, -\}$, let J_ε be the ideal in $\mathbb{A}$ generated by $f_\varepsilon, \rho^{-1} - \rho - (q^{-1} - q)(\delta_0 - 1)$, and the elements

$$\rho(\mathbf{a}_\ell - \mathbf{a}_{r-\ell}/\mathbf{a}_0) + (q - q^{-1}) \left[\sum_{j=1}^{r-\ell} \mathbf{a}_{j+\ell} \delta_j - \sum_{j=\max(\ell+1, \lceil r/2 \rceil)}^{\lfloor (\ell+r)/2 \rfloor} \mathbf{a}_{2j-\ell} + \sum_{j=\lceil \ell/2 \rceil}^{\min(\ell, \lceil r/2 \rceil - 1)} \mathbf{a}_{2j-\ell} \right],$$

for $1 \leq \ell \leq r-1$. Note that $J_0 \subseteq J_+ \cap J_-$. Set $R_\varepsilon = \mathbb{A}/J_\varepsilon$. Since $J_0 \subseteq J_\pm$, we have maps $R_0 \rightarrow R_\pm$ defined by $x + J_0 \mapsto x + J_\pm$. Moreover, we can define an automorphism θ of $\mathbb{A}$ taking J_+ to J_- . (For r odd, require $\theta(\rho) = -\rho, \theta(q) = -q$,

$\theta(\delta_j) = \delta_j$ and $\theta(u_i) = u_i$. For r even, require $\theta(\rho) = \rho$, $\theta(q) = -q^{-1}$, $\theta(\delta_j) = \delta_j$ and $\theta(u_i) = u_i$.) Then θ induces an isomorphism $\theta : R_+ \rightarrow R_-$.

The rings R_ε have the properties of an admissible ring, except that the image of $(q - q^{-1})$ in R_ε might be a zero-divisor. Consequently, we will introduce quotients of R_ε in which the image of $(q - q^{-1})$ is not a zero divisor.

If S is an admissible integral domain, then there is a morphism of ground rings from R_0 to S , and this homomorphism must factor through R_+ or R_- according to Remark 3.7. In particular, there is a morphism of ground rings from R_0 to the universal u -admissible ring $\overline{S}$ that factors through R_+ . It follows from this that $(q - q^{-1})^n + J_0 \neq 0$ in R_0 and $(q - q^{-1})^n + J_+ \neq 0$ in R_+ , for all natural numbers n . Using the isomorphism $\theta : R_+ \rightarrow R_-$, we have also that $(q - q^{-1})^n + J_- \neq 0$ in R_- for all natural numbers n .

For $\varepsilon \in \{0, +, -\}$, let

$$P_\varepsilon = \{x + J_\varepsilon \in R_\varepsilon : \exists n \geq 1 \text{ such that } (x + J_\varepsilon)(q - q^{-1} + J_\varepsilon)^n = 0\}.$$

Then P_ε is an ideal in R_ε . Set $\overline{R}_\varepsilon = R_\varepsilon/P_\varepsilon$. Let $\bar{\rho}$, $\bar{q}$, etc. denote the images of the parameters in $\overline{R}_\varepsilon$. Since the quotient maps $R_0 \rightarrow R_\pm$ take P_0 into $P_\pm$, they induce maps $\overline{R}_0 \rightarrow \overline{R}_\pm$. Moreover, the isomorphism $\theta : R_+ \rightarrow R_-$ maps P_+ to P_- , so induces an isomorphism $\theta : \overline{R}_+ \rightarrow \overline{R}_-$.

If S is any admissible ring and $\varphi : R_\varepsilon \rightarrow S$ is a morphism of ground rings from R_ε to S , then $\varphi(P_\varepsilon) = 0$. Hence φ induces a morphism of ground rings from $\overline{R}_\varepsilon$ to S . In particular, we have a morphism of ground rings from R_0 to the universal u -admissible ring $\overline{S}$, that factors through R_+ , and this map induces a morphism of ground rings from $\overline{R}_0$ to $\overline{S}$, factoring through $\overline{R}_+$. It follows from this that $\bar{q} - \bar{q}^{-1} \neq 0$ in $\overline{R}_\varepsilon$ ($\varepsilon = 0, +$). Using the isomorphism $\theta : \overline{R}_+ \rightarrow \overline{R}_-$, we see that $\bar{q} - \bar{q}^{-1} \neq 0$ in $\overline{R}_-$ as well.

Finally we check that $\bar{q} - \bar{q}^{-1}$ is not a zero-divisor in $\overline{R}_\varepsilon$, for $\varepsilon \in \{0, +, -\}$. Let $\bar{x} \in \overline{R}$ such that $\bar{x}(\bar{q} - \bar{q}^{-1}) = 0$. Let x be a preimage of $\bar{x}$ in R_ε . Then $x(q - q^{-1} + J_\varepsilon) \in P_\varepsilon$. Hence there exists an $n \geq 1$ such that

$$x((q - q^{-1}) + J_\varepsilon)((q - q^{-1}) + J_\varepsilon)^n = 0.$$

But this means that $x \in P_\varepsilon$, so $\bar{x} = 0$.

We claim that $\overline{R}_+$ is an integral domain.

Since there is a morphism of ground rings from $\overline{R}_+$ to the universal u -admissible ring $\overline{S}$, it follows that the parameters $\bar{q}, \bar{u}_1, \dots, \bar{u}_r$ in $\overline{R}_+$ are algebraically independent over $\mathbb{Z}$. Consider $\overline{R}_+[(\bar{q} - \bar{q}^{-1})^{-1}]$ and its subring

$$A_0 = \mathbb{Z}[\bar{q}^{\pm 1}, \bar{u}_1^{\pm 1}, \dots, \bar{u}_r^{\pm 1}][(\bar{q} - \bar{q}^{-1})^{-1}].$$

A_0 is an integral domain, since $\bar{q}, \bar{u}_1, \dots, \bar{u}_r$ are algebraically independent over $\mathbb{Z}$. According to Corollary 3.8, A_0 also contains $\bar{\rho}^{\pm 1}$ and $\bar{\delta}_0, \bar{\delta}_1, \dots, \bar{\delta}_{r-1}$. Therefore,

$$A_0[\bar{\delta}_0^{-1}] = \overline{R}_+[(\bar{q} - \bar{q}^{-1})^{-1}].$$

Thus, $\overline{R}_+[(\bar{q} - \bar{q}^{-1})^{-1}]$ and its subring $\overline{R}_+$ are integral domains. According to Corollary 3.12, $\overline{R}_+$ is u -admissible. It is now clear that $\overline{R}_+[(\bar{q} - \bar{q}^{-1})^{-1}]$ is isomorphic to the universal u -admissible ring $\overline{S}$ by a isomorphism of ground rings.

We summarize this discussion with the following theorem:

Theorem 3.19. *There exist a universal admissible ring $\overline{R}_0$ and a universal admissible integral domain $\overline{R}_+$ with the following properties:*

- (1) *There is morphism of ground rings from $\overline{R}_0$ to $\overline{R}_+$.*
- (2) *$\overline{R}_0$ (and hence also $\overline{R}_+$) are generated as unital rings by the parameters $\bar{q}^{\pm 1}, \bar{\rho}^{\pm 1}, \bar{\delta}_0^{\pm 1}, \bar{\delta}_1, \dots, \bar{\delta}_{r-1}, \bar{u}_1^{\pm 1}, \dots, \bar{u}_r^{\pm 1}$.*
- (3) *The parameters $\bar{q}, \bar{u}_1, \dots, \bar{u}_r$ of $\overline{R}_+$ (hence also of $\overline{R}_0$) are algebraically independent over $\mathbb{Z}$.*
- (4) *Whenever S is a ring with admissible parameters, there exists a morphism of ground rings from $\overline{R}_0$ to S . If S is also an integral domain, then the morphism of ground rings can be chosen to factor through $\overline{R}_+$.*
- (5) *$\overline{R}_+[(\bar{q} - \bar{q}^{-1})^{-1}]$ is isomorphic to the universal u -admissible ring, by an isomorphism of ground rings.*

3.5. More generating functions. Let F be a field with u -admissible parameters $\rho, q, \delta_j, u_1, \dots, u_r$. Let $KT_{n,r}$ denote $KT_{n,F,r}(u_1, \dots, u_r)$. Recall that ε_n denotes the conditional expectation from $KT_{n,r}$ to $KT_{n-1,r}$, and ε denotes the canonical trace on $KT_{n,r}$. For $n \geq 1$, write Y_n for $\varphi(y_n)$. Note that $\varepsilon_n(Y_n^a) \in Z(KT_{n-1,r})$ for $a \geq 0$, since Y_n^a commutes with $KT_{n-1,r}$. Define $\omega_n^{(a)}$ by $\omega_n^{(a)} = \delta_0 \varepsilon_n(Y_n^a)$, so

$$E_n Y_n^a E_n = \omega_n^{(a)} E_n.$$

Define

$$(3.25) \quad Z_n(t) = \sum_{a \geq 0} \omega_n^{(a)} t^{-a},$$

a formal power series with coefficients in the center of $KT_{n-1,r}$. This is consistent with the previous definition of Z_1 in 3.23. Moreover, define

$$(3.26) \quad Q_n(t) = Z_n(t) - \rho^{-1}/(q^{-1} - q) - t^2/(t^2 - 1).$$

In particular

$$(3.27) \quad Q_1(t) = G(t^{-1})A(t),$$

according to (3.24).

We have the following remarkable recursion for $Q_n(t)$, which is due to Beliakova and Blanchet [5], Lemma 7.4.

Proposition 3.20.

$$Q_{n+1}(t) = Q_n(t) \left(\frac{(t - Y_n)^2(t - q^{-2}Y_n^{-1})(t - q^2Y_n^{-1})}{(t - Y_n^{-1})^2(t - q^{-2}Y_n)(t - q^2Y_n)} \right).$$

Proof. Beliakova and Blanchet prove this for the ordinary BMW (or Kauffman tangle) algebras, that is, for $r = 1$. However, their proof remains valid in the general case. $\square$

4. THE GENERIC STRUCTURE OF THE CYCLOTOMIC BMW ALGEBRA

In this section we obtain the “generic structure” of the cyclotomic BMW algebras over a field, following the method of Wenzl from [6], [29], and [30]. The argument also relies on ideas from Beliakova and Blanchet [5].

Let F be the field of fractions of the universal admissible integral domain $\overline{R}_+$. We will show that for all n the cyclotomic BMW algebra $W_{n,F,r}$ over F is isomorphic to the cyclotomic Kauffman tangle algebra $\widehat{KT}_{n,F,r}$ over F , and is split semisimple of dimension $r^n(2n - 1)!!$. The irreducible representations of $W_{n,F,r}$ are labelled by r -tuples of Young diagrams of total size $n, n - 2, n - 4, \dots$, and the dimension of the irreducible representation labelled by such an r -tuple μ is the number of up-down tableaux of length n and shape μ .

4.1. Semisimplicity and structure of $W_{n,F,r}$. Let S be a ground ring. For each n , let I_n be the two sided ideal in $W_{n,S,r}$ generated by e_{n-1} . Because of the relations $e_j e_{j\pm 1} e_j = e_j$, the ideal I_n is generated by any e_i ($1 \leq i \leq n - 1$) or by all of them.

Lemma 4.1. *For all $n \geq 1$, $I_{n+1} = W_{n,S,r} e_n W_{n,S,r}$.*

Proof. This follows from the corresponding result for the affine BMW algebras, [9], Proposition 3.20. $\square$

Let us recall the definition of the affine and cyclotomic Hecke algebras, see [1].

Definition 4.2. Let S be a commutative unital ring with an invertible element q . The *affine Hecke algebra* $\hat{H}_{n,S}(q^2)$ over S is the S -algebra with generators $t_1, g_1, \dots, g_{n-1}$, with relations:

- (1) The generators g_i are invertible, satisfy the braid relations, and $g_i - g_i^{-1} = (q - q^{-1})$.
- (2) The generator t_1 is invertible, $t_1 g_1 t_1 g_1 = g_1 t_1 g_1 t_1$ and t_1 commutes with g_j for $j \geq 2$.

Let $u_1, \dots, u_r$ be additional elements in S . The *cyclotomic Hecke algebra* $H_{n,S,r}(q^2; u_1, \dots, u_r)$ is the quotient of the affine Hecke algebra $\hat{H}_{n,S}(q^2)$ by the polynomial relation $(t_1 - u_1) \cdots (t_1 - u_r) = 0$.

Remark 4.3. Since the generator t_1 can be rescaled by an arbitrary invertible element of S , only the ratios of the parameters u_i have invariant significance in the definition of the cyclotomic Hecke algebra.

Lemma 4.4. Let S be a ground ring with parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. For all $n \geq 1$, the quotient of the cyclotomic BMW algebra $W_{n,S,r}(u_1, \dots, u_r)$ by the ideal I_n is isomorphic to the cyclotomic Hecke algebra $H_{n,S,r}(q^2; u_1, \dots, u_r)$.

Proof. Evident. $\square$

Lemma 4.5. Let F be a field with u -admissible parameters ρ, q, δ_j ($j \geq 0$), and $u_1, \dots, u_r$. Assume that $u_j \notin \{\pm q^{\pm 1}\}$ for all j .

- (1) $W_{1,F,r} \cong KT_{1,F,r} \cong F^r$.
- (2) Let $P_j = \prod_{\ell \neq j} \frac{Y_1 - u_\ell}{u_j - u_\ell} \in KT_{1,F,r}$. Then $EP_j E = \gamma_j E$, and $\varepsilon(P_j) = \gamma_j / \delta_0$, where γ_j is defined by (3.6).

Proof. Let $\varphi : W_{2,F,r} \rightarrow KT_{2,F,r}$ be the surjective algebra homomorphism determined by $\varphi(y_1) = Y_1$, $\varphi(e) = E$, $\varphi(g) = G$. Define $p_j = \prod_{\ell \neq j} \frac{y_1 - u_\ell}{u_j - u_\ell} \in W_{1,F,r}$. It is shown in the proof of Theorem 3.10 that $ep_j e = \gamma_j e$. Hence, $EP_j E = \varphi(ep_j e) = \gamma_j E$.

By Remark 3.14, $\gamma_j \neq 0$ for all j , and by Lemma 2.16, $E \neq 0$. Therefore, $P_j \neq 0$. The elements P_j are mutually orthogonal non-zero idempotents, so they are linearly independent. Thus the restriction of φ to $W_{1,F,r}$ is an isomorphism. Moreover, $KT_{1,F,r} \cong FP_1 \oplus \cdots \oplus FP_r \cong F^r$, as algebras. $\square$

Let $\lambda = (\lambda^{(1)}, \dots, \lambda^{(r)})$ be an r -tuple of Young diagrams. The total size of λ is $|\lambda| = \sum_i |\lambda^{(i)}|$. If μ and λ are r -tuples of Young diagrams of total size $f - 1$ and f respectively, we write $\mu \subset \lambda$ if μ is obtained from λ by removing one box from one component of λ .

Theorem 4.6 ([1]). *Let F be a field. The cyclotomic Hecke algebra $H_{n,F,r}(q; u_1, \dots, u_r)$ is split semisimple for all n as long as q is not a proper root of unity and u_i/u_j is not a power of q for all $i \neq j$. In this case, the simple components of $H_{n,F,r}(q; u_1, \dots, u_r)$ are labeled by r -tuples of Young diagrams of total size n , and a simple $H_{n,F,r}$ module V_λ decomposes as a $H_{n-1,F,r}$ module as the direct sum of all V_μ with $\mu \subset \lambda$.*

In the following, let Γ_n denote the set of r -tuples of Young diagrams whose total size is no more than n and congruent to $n \bmod 2$. If $\lambda \in \Gamma_n$ and $\mu \in \Gamma_{n-1}$, write $\mu \leftrightarrow \lambda$ if μ is obtained from λ by either adding one box to, or removing one box from, one component of λ . For $\lambda \in \Gamma_n$, an *up-down tableau of length n and shape λ* is a sequence $(\lambda^{(0)}, \lambda^{(1)}, \dots, \lambda^{(n)})$, where

- (1) for all i , $\lambda^{(i)} \in \Gamma_i$,
- (2) $\lambda^{(0)} = (\emptyset, \dots, \emptyset)$,
- (3) $\lambda^{(n)} = \lambda$, and
- (4) for each i , $\lambda^{(i)} \leftrightarrow \lambda^{(i+1)}$.

Let $\mathcal{T}(n)$ denote the set of up-down tableaux of length n , and $\mathcal{T}(n, \lambda)$ the set of up-down tableaux of length n and shape λ .

Definition 4.7. Let λ be an r -tuple of Young diagrams. Label the nodes (or cells, or boxes) of λ by triples (j, x, y) , where j is the component ($1 \leq j \leq r$), x is the row coordinate, and y is the column coordinate of the node. Define the *multiplicative content* of a node $\alpha = (j, x, y)$ by $\tilde{c}(\alpha) = u_j q^{2(y-x)}$.

An *addable node* of λ is a node which can be added to one component to give a new r -tuple of Young diagrams. A *removable node* is a node which can be removed with the result still being an r -tuple of Young diagrams.

For an addable node of λ , set $b(\alpha, \lambda) = \tilde{c}(\alpha)$, and for removable node, set $b(\alpha, \lambda) = \tilde{c}(\alpha)^{-1}$.

If $T = (\lambda^{(0)}, \lambda^{(1)}, \dots, \lambda^{(n)})$ is an up-down tableau of length n and $j \leq n$, let $b(j, T) = b(\alpha_j, \lambda^{(j-1)})$, where α_j is the node that is added or removed in passing from $\lambda^{(j-1)}$ to $\lambda^{(j)}$.

The total number of addable and removable nodes of a single Young diagram is always odd, so the the total number of addable and removable nodes of an r -tuple of Young diagrams has the same parity as r .

In the following we work over the field of fractions F of the universal admissible integral domain $\overline{R}_+$. Since $\overline{R}_+$ imbeds in the universal u -admissible ring, according to Theorem 3.19, we will now denote the parameters of $\overline{R}_+$ by ρ , q , δ_j , and u_i . Recall that $q, u_1, \dots, u_r$ are algebraically independent over $\mathbb{Z}$.

Theorem 4.8. *Let F denote the field of fractions of the universal admissible integral domain $\overline{R}_+$. Write W_n for $W_{n,F,r}(u_1, \dots, u_r)$, KT_n for $KT_{n,F,r}(u_1, \dots, u_r)$, and H_n for $H_{n,F,r}(q^2; u_1, \dots, u_r)$. The following statements hold for all $n \geq 0$.*

- (1) $\varphi : W_n \rightarrow KT_n$ is an isomorphism.
- (2) The Markov trace ε on KT_n is non-degenerate.
- (3) W_n is split semisimple, and $W_n \cong I_n \oplus H_n$.
- (4) The minimal ideals of W_n are labeled by Γ_n ; more precisely, the minimal ideals of I_n are labeled by $\{\lambda \in \Gamma_n : |\lambda| < n\}$ and the minimal ideals in W_n/I_n are labeled by $\{\lambda \in \Gamma_n : |\lambda| = n\}$.
- (5) The branching diagram for $W_{n-1} \subseteq W_n$ has a single edge connecting $\mu \in \Gamma_{n-1}$ and $\lambda \in \Gamma_n$ if $\mu \leftrightarrow \lambda$, and no edges otherwise.
- (6) The set of paths on the branching diagram for the sequence $W_0 \subseteq W_1 \subseteq \dots \subseteq W_n$ is the set of up-down tableaux of length n .
- (7) Let p_T be the path idempotent in W_n corresponding to an up-down tableau T of length n . Then p_T is a common eigenvector for $y_1, y_2, \dots, y_n$,

$$y_k p_T = p_T y_k = b(k, T) p_T.$$

Proof. The assertions of the theorem hold trivially for $n = 0$, and the assertions for $n = 1$ are implied by Lemma 4.5. (We label the one minimal ideal of $W_{0,F} \cong F$ by the r -tuple of empty diagrams, $\emptyset = (\emptyset, \dots, \emptyset)$. We label the minimal ideal Fp_j of W_1 by the the r -tuple $\square_j$, with the Young diagram $\square$ in the j -th position, and the empty diagram $\emptyset$ in all other positions. Assertion (2) holds for $n = 1$ since

$\varepsilon(p_j) = \gamma_j/\delta_0$ by Lemma 4.5 and $\gamma_j \neq 0$ by Remark 3.14. Assertion (7) holds for $n = 1$ because $y_1 p_j = u_j p_j$, and u_j is the multiplicative content of the node added to $(\emptyset, \dots, \emptyset)$ to get $\square_j$.)

We proceed by induction on n . Fix $n \geq 1$, and suppose that the assertions hold for W_f for $f \leq n$.

We have a conditional expectation $\varepsilon_n : W_n \rightarrow W_{n-1}$ that preserves the non-degenerate trace on W_n and W_{n-1} ; that is, $\varepsilon \circ \varepsilon_n = \varepsilon$. (The conditional expectation results from the isomorphism $W_f \cong KT_f$ for $f \leq n$ and the conditional expectation on the cyclotomic Kauffman tangle algebras; see Section 2.6.) Moreover, the conditional expectation is implemented by the idempotent $e = (1/\delta_0)e_n \in W_{n+1}$; that is $ebe = \varepsilon_n(b)e$ for $b \in W_n$. Therefore, by Wenzl's Theorem 2.17 and the induction hypothesis, $W_n e_n W_n \subseteq W_{n+1}$ is split semisimple, with minimal ideals labeled by Γ_{n-1} . By Lemma 4.1, $W_n e_n W_n = I_{n+1}$.

Since $I_{n+1} = W_n e_n W_n$ has an identity element z , necessarily a central idempotent in W_{n+1} , it follows that I_{n+1} has a complementary ideal J_{n+1} ; moreover, $J_{n+1} \cong H_{n+1}$, according to Lemma 4.4. By Theorem 4.6 and the algebraic independence of $\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r$ over $\mathbb{Z}$, H_{n+1} is split semisimple. Thus W_{n+1} is split semisimple and $W_{n+1} \cong I_{n+1} \oplus H_{n+1}$. This proves statement (3) for W_{n+1} .

To prove statements (4), (5) and (6) for W_{n+1} , we have to determine the branching diagram for $W_n \subseteq W_{n+1}$. We know that the multiplicity matrix for the map $x \mapsto xz$, from W_n to I_{n+1} , is the transpose of the inclusion matrix for $W_{n-1} \subseteq W_n$, by Theorem 2.17. The map $x \mapsto x(1-z)$, from I_n to J_{n+1} , is zero, since $I_n \subseteq I_{n+1}$. The multiplicity matrix for the map $x \mapsto x(1-z)$, from J_n to J_{n+1} is the same as that for the inclusion $H_n \subseteq H_{n+1}$, which is given by Theorem 4.6. Statements (4), (5), and (6) follow from these observations.

It remains to prove statements (1), (2) and (7) for W_{n+1} and KT_{n+1} .

Recall that for any up-down tableau T of length $k \leq n+1$, T' denotes the truncation of T of length $k-1$. If T is an up-down tableau of length k and shape μ , and if $\lambda \in \Gamma_{k+1}$ satisfies $\mu \leftrightarrow \lambda$, then $T + \lambda$ denotes the extension of T of length $k+1$ and shape λ . Let p_T denote the path idempotent in W_k corresponding to an up-down tableau T of length k , and let $P_T = \varphi(p_T)$ be its image in KT_k .

Lemma 4.9. *Assume that assertions (1), (2), and (7) hold for W_k and KT_k for all $k \leq n$. If T is an up-down tableau of length $n+1$, then for all $k \leq n+1$,*

$$y_k p_T = p_T y_k = b(k, T) p_T.$$

That is, assertion (7) also holds for W_{n+1} .

Proof of Lemma 4.9. Let T be an up-down tableau of length $n+1$. If $k \leq n$, then

$$y_k p_T = y_k p_{T'} p_T = b(k, T') p_{T'} p_T = b(k, T) p_T,$$

and similarly for $p_T y_k$.

It remains to show that $y_{n+1} p_T = p_T y_{n+1} = b(n+1, T) p_T$.

Write $T = (\emptyset, \lambda^{(1)}, \dots, \lambda^{(n-1)}, \mu, \lambda)$. In case $|\lambda| = n+1$, p_T belongs to the ideal J_{n+1} , which is isomorphic to the cyclotomic Hecke algebra H_{n+1} . The result then follows from formulas for the irreducible representations of the cyclotomic Hecke algebras, see [2, 1].

We assume now that $|\lambda| \leq n-1$. Consider the minimal ideal $W_{n,\mu}$ of W_n corresponding to μ . This ideal is isomorphic to a full matrix algebra over F , and the path idempotents p_U ($U \in \mathcal{T}(n, \mu)$) are a complete family of mutually orthogonal minimal idempotents in $W_{n,\mu}$. Hence $W_{n,\mu}$ has a system of matrix units $e_{S,U}$ ($S, U \in \mathcal{T}(n, \mu)$) such that $p_U = e_{U,S} p_S e_{S,U}$. Thus for any $S \in \mathcal{T}(n, \mu)$, we can write

$$\begin{aligned} p_T &= p_{T'} z_{\lambda}^{(n+1)} = e_{T',S} p_S e_{S,T'} z_{\lambda}^{(n+1)} \\ &= e_{T',S} p_S z_{\lambda}^{(n+1)} e_{S,T'} = e_{T',S} p_{S+\lambda} e_{S,T'}. \end{aligned}$$

Suppose that there exists some $S \in \mathcal{T}(n, \mu)$ such that

$$y_{n+1} p_{S+\lambda} = p_{S+\lambda} y_{n+1} = b(\alpha, \mu) p_{S+\lambda},$$

where α is the node added or removed in passing from μ to λ . Then we have

$$\begin{aligned} y_{n+1} p_T &= y_{n+1} e_{T',S} p_{S+\lambda} e_{S,T'} = e_{T',S} y_{n+1} p_{S+\lambda} e_{S,T'} \\ &= b(\alpha, \mu) e_{T',S} p_{S+\lambda} e_{S,T'} = b(\alpha, \mu) p_T, \end{aligned}$$

and similarly for $p_T y_{n+1}$. Therefore, it suffices to prove the result when T has the form $T = (\emptyset, \lambda^{(1)}, \dots, \lambda, \mu, \lambda)$. Let T' be the truncation of T of length n and T'' the truncation of length $n-1$. Then $p_{T''}$ is a minimal idempotent in the minimal ideal $W_{n-1,\lambda}$ of W_{n-1} . Let $e = (1/\delta_0) e_n$, so that $ebe = \varepsilon_n(b)e$ for $b \in W_n$. Then by Remark 2.18, $p_{T''} e$ is a minimal idempotent in $W_{n+1,\lambda}$.

Consider the element $p_{T'} e p_{T'}$. We have

$$\begin{aligned} (p_{T'} e p_{T'})^2 &= p_{T'} e \varepsilon_n(p_{T'}) p_{T'} = (\varepsilon(p_{T'})/\varepsilon(p_{T''})) p_{T'} e p_{T''} p_{T'} \\ &= (\varepsilon(p_{T'})/\varepsilon(p_{T''})) p_{T'} e p_{T'}, \end{aligned}$$

using Lemma 2.19, so $p_{T'} e p_{T'}$ is an essential idempotent. Write $\kappa = \varepsilon(p_{T'})/\varepsilon(p_{T''})$ and $f = \kappa^{-1} p_{T'} e p_{T'}$. (We are using the assumption that the trace is non-degenerate on W_n and W_{n-1} , which follows from assertions (1) and (2) for these algebras.)

We have $f = \kappa^{-1} p_{T'} (p_{T''} e) p_{T'}$, which shows that f is in $W_{n+1, \lambda}$. But then $f = z_{\lambda}^{(n+1)} f z_{\lambda}^{(n+1)} = (z_{\lambda}^{(n+1)} p_{T'}) f (p_{T'} z_{\lambda}^{(n+1)}) = p_T f p_T$. Therefore f is a multiple of, and hence equal to, the minimal idempotent p_T . Now we can compute:

$$\begin{aligned} y_{n+1} p_T &= \kappa^{-1} y_{n+1} p_{T'} e p_{T'} = \kappa^{-1} p_{T'} y_{n+1} e p_{T'} \\ &= \kappa^{-1} p_{T'} y_n^{-1} e p_{T'} = \kappa^{-1} y_n^{-1} p_{T'} e p_{T'} \\ &= y_n^{-1} p_T = b(\alpha, \mu) p_T, \end{aligned}$$

and similarly for $p_T y_{n+1}$. This completes the proof of Lemma 4.9.

Recall the quantities $Z_{k+1}(t)$ and $Q_{k+1}(t)$ from Section 3.5. They are formal power series with coefficients in the center of KT_k . We want to find an expression for $Q_{k+1}(t) p_T$, where T is an up-down tableau of length k .

Let λ be any r -tuple of Young diagrams. Define

$$(4.1) \quad \tilde{Q}(t, \lambda) = p A(t) \prod_{\alpha} \frac{t - b(\alpha, \lambda)^{-1}}{t - b(\alpha, \lambda)},$$

where $p = \prod_j u_j$, $A(t)$ is as in (3.24), and α runs over all addable and removable nodes of λ . Define

$$(4.2) \quad \tilde{Z}(t, \lambda) = \tilde{Q}(t, \lambda) + \rho^{-1}/(q^{-1} - q) + t^2/(t^2 - 1).$$

Note that if $\emptyset$ is the r -tuple of empty Young diagrams, then

$$(4.3) \quad \tilde{Q}(t, \emptyset) = p A(t) \prod_j \frac{t - u_j^{-1}}{t - u_j} = A(t) G(t^{-1}) = Q_1(t),$$

using (3.27) and thus $\tilde{Z}(t, \emptyset) = Z_1(t)$.

Lemma 4.10. *Suppose μ and λ are r -tuples of Young diagrams and λ is obtained from μ by adding a single node α . Let $\tilde{c} = \tilde{c}(\alpha)$ be the multiplicative content of this node. Then*

$$\tilde{Q}(t, \lambda) = \tilde{Q}(t, \mu) \frac{(t - \tilde{c})^2 (t - q^{-2} \tilde{c}^{-1}) (t - q^2 \tilde{c}^{-1})}{(t - \tilde{c}^{-1})^2 (t - q^{-2} \tilde{c}) (t - q^2 \tilde{c})}$$

Proof of Lemma 4.10. Suppose $\alpha = (j, x, y)$, so $y - 1 = \mu_x^{(j)}$, the length of the x -th row of the j -th component of $\boldsymbol{\mu}$. There are several cases to consider according to the relative size of $\mu_{x-1}^{(j)}$, $\mu_x^{(j)}$, and $\mu_{x+1}^{(j)}$. For example, suppose $\mu_x^{(j)} = \mu_{x+1}^{(j)}$, and $\mu_x^{(j)} + 1 < \mu_{x-1}^{(j)}$. Then adding the node α :

- (1) eliminates the addable node at α and produces a removable node at α , and
- (2) produces two new addable nodes at $(j, x, y + 1)$ and $(j, x + 1, y)$.

The node at α contributes $((t - \tilde{c})/(t - \tilde{c}^{-1}))^2$ to the ratio $\tilde{Q}(t, \boldsymbol{\lambda})/\tilde{Q}(t, \boldsymbol{\mu})$. The two new addable nodes contribute

$$\left(\frac{t - q^{-2}\tilde{c}^{-1}}{t - q^2\tilde{c}} \right) \left(\frac{t - q^2\tilde{c}^{-1}}{t - q^{-2}\tilde{c}} \right).$$

Other cases can be analyzed similarly. This completes the proof of Lemma 4.10.

Lemma 4.11. *Let T be an up-down tableau of length $k \leq n + 1$ and shape $\boldsymbol{\lambda}$. Then*

$$Q_{k+1}(t)P_T = \tilde{Q}(t, \boldsymbol{\lambda})P_T.$$

and

$$Z_{k+1}(t)P_T = \tilde{Z}(t, \boldsymbol{\lambda})P_T.$$

Proof of Lemma 4.11. We do this by induction on k . There is a unique tableau of length 0, and the corresponding idempotent is the identity. So the statement for $k = 0$ reads $Q_1(t) = \tilde{Q}(t, \emptyset)$, which was observed in (4.3). Let T be an up-down tableau of length $k \geq 1$ and suppose the assertion holds for all shorter up-down tableaux. Let $\boldsymbol{\lambda}$ denote the shape of T and $\boldsymbol{\mu}$ the shape of T' . Then

$$\begin{aligned} Q_{k+1}(t)P_T &= Q_k(t) \left(\frac{(t - Y_k)^2(t - q^{-2}Y_k^{-1})(t - q^2Y_k^{-1})}{(t - Y_k^{-1})^2(t - q^{-2}Y_k)(t - q^2Y_k)} \right) P_{T'}P_T \\ &= Q_k(t)P_{T'} \left(\frac{(t - Y_k)^2(t - q^{-2}Y_k^{-1})(t - q^2Y_k^{-1})}{(t - Y_k^{-1})^2(t - q^{-2}Y_k)(t - q^2Y_k)} \right) P_T. \end{aligned}$$

By the induction hypothesis and Lemma 4.9, this is equal to

$$\begin{aligned} \tilde{Q}(t, \boldsymbol{\mu})P_{T'} &\left(\frac{(t - b)^2(t - q^{-2}b^{-1})(t - q^2b^{-1})}{(t - b^{-1})^2(t - q^{-2}b)(t - q^2b)} \right) P_T \\ &= \tilde{Q}(t, \boldsymbol{\mu}) \left(\frac{(t - b)^2(t - q^{-2}b^{-1})(t - q^2b^{-1})}{(t - b^{-1})^2(t - q^{-2}b)(t - q^2b)} \right) P_T, \end{aligned}$$

where $b = b(\alpha, \mu)$. By Lemma 4.10, this equals $\tilde{Q}(t, \lambda)P_T$. (In case $\lambda \subseteq \mu$, reverse the roles of λ and μ in the Lemma.) The conclusions of Lemma 4.11 follow from this.

Let U be an up-down tableau of length n . Note that P_U is a minimal idempotent in KT_n and $\varepsilon(P_U) \neq 0$ by claims (1) and (2) of the induction hypothesis. We have $P_U = \sum_T P_T$, where T ranges over all up-down tableaux of length $n+1$ such that $T' = U$. It follows that there exists some T such that $\varepsilon(P_T) \neq 0$. Fix such a T .

Recall from Lemma 2.19 that for any up-down tableau S of length $n+1$ such that $S' = U$, we have

$$(4.4) \quad \varepsilon_{n+1}(P_S) = (\varepsilon(P_S)/\varepsilon(P_U))P_U.$$

Next, for any such up-down tableau S , we claim

$$(4.5) \quad \varepsilon(P_T E_{n+1} P_S E_{n+1} P_T) = \frac{\varepsilon(P_T) \varepsilon(P_S)}{\varepsilon(P_U)}.$$

In fact,

$$\begin{aligned} \varepsilon(P_T E_{n+1} P_S E_{n+1} P_T) &= \varepsilon(E_{n+1} P_S E_{n+1} P_T) \\ &= \delta_0 \varepsilon(E_{n+1} \varepsilon_{n+1}(P_S) P_T) = \delta_0 \frac{\varepsilon(P_S)}{\varepsilon(P_U)} \varepsilon(E_{n+1} P_{T'} P_T) \\ &= \delta_0 \frac{\varepsilon(P_S)}{\varepsilon(P_U)} \varepsilon(E_{n+1} P_T) = \frac{\varepsilon(P_T) \varepsilon(P_S)}{\varepsilon(P_U)}, \end{aligned}$$

where we have used Lemma 2.15 and (4.4).

Now consider the formal power series:

$$(4.6) \quad \begin{aligned} E_{n+1}(t - Y_{n+1})^{-1} E_{n+1} &= E_{n+1}(1/t) \sum_{a \geq 0} Y_{n+1}^a t^{-a} E_{n+1} \\ &= (1/t) Z_{n+1}(t) E_{n+1} \end{aligned}$$

Let U and T be as above, so $T' = U$ and $\varepsilon(P_T) \neq 0$. Denote the shape of U by μ . Multiply both sides of 4.6 by P_T , and use Lemma 4.11 to get

$$(4.7) \quad P_T E_{n+1}(t - Y_{n+1})^{-1} E_{n+1} P_T = (1/t) \tilde{Z}(t, \mu) P_T E_{n+1} P_T.$$

Now take the trace on both sides,

$$(4.8) \quad \varepsilon(P_T E_{n+1}(t - Y_{n+1})^{-1} E_{n+1} P_T) = (1/t) \tilde{Z}(t, \mu) \delta_0^{-1} \varepsilon(P_T).$$

Now we focus on the left side of the equation. Note that for S an up-down tableau of length $n+1$, $P_S E_{n+1} P_T = P_S P_{S'} P_{T'} E_{n+1} P_T$, and this is non-zero

only if $S' = T' = U$. Therefore, the left side equals

$$\sum_S \varepsilon(P_T E_{n+1} (t - Y_{n+1})^{-1} P_S E_{n+1} P_T),$$

where the sum is over all S such that $S' = U$. By Lemma 4.9, and (4.5) this equals

$$\begin{aligned} & \sum_S (t - b(n+1, S))^{-1} \varepsilon(P_T E_{n+1} P_S E_{n+1} P_T) \\ (4.9) \quad &= \sum_S (t - b(n+1, S))^{-1} \frac{\varepsilon(P_S) \varepsilon(P_T)}{\varepsilon(P_U)} \\ &= \sum_{\alpha} (t - b(\alpha, \boldsymbol{\mu}))^{-1} \frac{\varepsilon(P_{S(\alpha)}) \varepsilon(P_T)}{\varepsilon(P_U)}, \end{aligned}$$

where the last sum is over all addable and removable nodes of $\boldsymbol{\mu}$, and $S(\alpha)$ is the extension of U by α ; that is, if $U = (\boldsymbol{\lambda}^{(0)}, \dots, \boldsymbol{\mu})$, then $S(\alpha) = (\boldsymbol{\lambda}^{(0)}, \dots, \boldsymbol{\mu}, \boldsymbol{\lambda}(\alpha))$, where $\boldsymbol{\lambda}(\alpha)$ is obtained from $\boldsymbol{\mu}$ by addition or removal of the node α . Comparing 4.8 and 4.9 and cancelling $\varepsilon(P_T)$, we have

$$(4.10) \quad (1/t) \tilde{Z}(t, \boldsymbol{\mu}) = \delta_0 \sum_{\alpha} (t - b(\alpha, \boldsymbol{\mu}))^{-1} \frac{\varepsilon(P_{S(\alpha)})}{\varepsilon(P_U)},$$

Since $\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r$ are algebraically independent over $\mathbb{Z}$, if α and β are two different addable or removable nodes of $\boldsymbol{\mu}$, then $\tilde{c}(\alpha) \neq \tilde{c}(\beta)^{\pm 1}$; hence $b(\alpha, \boldsymbol{\mu}) \neq b(\beta, \boldsymbol{\mu})$. Let S be any particular extension of U , $S = S(\beta)$, where β is an addable or removable node of $\boldsymbol{\mu}$; let $b = b(\beta, \boldsymbol{\mu})$. Take the residue at $t = b$ on both sides of 4.10. On the left side, we get

$$\begin{aligned} & \text{Res}(1/t) \tilde{Z}(t, \boldsymbol{\mu})|_{t=b} = \text{Res}(1/t) \tilde{Q}(t, \boldsymbol{\mu})|_{t=b} \\ (4.11) \quad &= p b^{-1} A(b) (b - b^{-1}) \prod_{\alpha \neq \beta} \frac{b - b(\alpha, \boldsymbol{\mu})^{-1}}{b - b(\alpha, \boldsymbol{\mu})}, \end{aligned}$$

while on the right side we get $\delta_0 \varepsilon(P_S) / \varepsilon(P_U)$. Therefore,

$$(4.12) \quad \varepsilon(P_S) = \delta_0^{-1} p b^{-1} A(b) (b - b^{-1}) \prod_{\alpha \neq \beta} \frac{b - b(\alpha, \boldsymbol{\mu})^{-1}}{b - b(\alpha, \boldsymbol{\mu})} \varepsilon(P_U).$$

If r is odd, $A(b)(b - b^{-1}) = \frac{b - b^{-1}}{q - q^{-1}} + 1$, and $p = \boldsymbol{\rho}$; see (3.24). Therefore,

$$(4.13) \quad \varepsilon(P_S) = \delta_0^{-1} \boldsymbol{\rho} b^{-1} \left(\frac{b - b^{-1}}{q - q^{-1}} + 1 \right) \prod_{\alpha \neq \beta} \frac{b - b(\alpha, \boldsymbol{\mu})^{-1}}{b - b(\alpha, \boldsymbol{\mu})} \varepsilon(P_U).$$

If r is even, $A(b)(b - b^{-1}) = \frac{q^{-1}b - qb^{-1}}{q - q^{-1}}$, and $p = q^{-1}\rho$. Therefore,

$$(4.14) \quad \varepsilon(P_S) = \delta_0^{-1} \rho q^{-1} b^{-1} \left(\frac{q^{-1}b - qb^{-1}}{q - q^{-1}} \right) \prod_{\alpha \neq \beta} \frac{b - b(\alpha, \mu)^{-1}}{b - b(\alpha, \mu)} \varepsilon(P_U).$$

The algebraic independence of $\mathbf{q}, \mathbf{u}_1, \dots, \mathbf{u}_r$ over $\mathbb{Z}$ implies that $\varepsilon(P_S) \neq 0$.

The computation of $\varepsilon(P_S)$ given here was adapted from [5], Theorem 7.

We have shown that for every up-down tableau S of length $n + 1$, $\varepsilon(P_S) = \varepsilon(\varphi(p_S)) \neq 0$. Hence ε is a non-degenerate trace on W_{n+1} . Moreover, this shows that φ is not zero on any minimal ideal of W_{n+1} , and hence φ is injective on W_{n+1} . This proves points (1) and (2) for W_{n+1} , and completes the proof of the theorem. $\square$

Remark 4.12. (A semisimplicity criterion) Let K be a field with u -admissible parameters. The inductive proof of the theorem applies to $W_{n,K,r}$ as long as the weights of the Markov trace remain non-zero. For this, what is needed is that for all r -tuples of Young diagrams μ of size $\leq n$, and for all addable or removable nodes α and β of μ , with $b = b(\beta, \mu)$ and $b' = b(\alpha, \mu)$, we have

$$A(b)(b - b^{-1}) \neq 0, \text{ and } b \neq (b')^{\pm 1}.$$

For this, it suffices that for all j , $u_j \neq \pm q^{\pm(2k-1)}$, and for all j, j' , $u_j/u_{j'}$ and $u_j u_{j'}$ are not equal to q^{2k} for $k \leq n$. In particular, we have the following result:

Corollary 4.13. *Let K be a field with u -admissible parameters. Suppose that for all j, j' , none of the quantities $\pm u_j$, $u_j/u_{j'}$ and $u_j u_{j'}$ is equal to an integer power of q . Then $W_{n,K,r}$ is split semisimple for all n .*

5. THE $\mathbb{Z}_r$ -BRAUER ALGEBRA AND THE DIMENSION OF $W_{n,F,r}$.

Recall that for $\lambda \in \Gamma_n$, the set of up-down tableaux of length n and shape λ is denoted by $\mathcal{T}(n, \lambda)$. The dimension of the simple $W_{n,F,r}$ module labelled by $\lambda \in \Gamma_n$ is $|\mathcal{T}(n, \lambda)|$, and the dimension of $W_{n,F,r}$ over F is the $\sum_{\lambda \in \Gamma_n} |\mathcal{T}(n, \lambda)|^2$. It should be possible to compute this quantity by an appropriate Shensted type correspondence, but it is more convenient to exhibit another sequence of split semisimple algebras $D_0 \subseteq D_1 \subseteq \dots \subseteq D_n$ (over some field) whose dimensions are known and which have the same branching diagram.

For this purpose, we introduce the $\mathbb{Z}_r$ -Brauer algebras. The $\mathbb{Z}_r$ - and $\mathbb{Z}$ -Brauer algebras were considered in [12], [27], and [9]. The G -Brauer algebras for an arbitrary abelian group G were studied in [25].

We recall that a Brauer diagram is a tangle diagram in the plane, in which information about over- and under-crossings is ignored: An (n, n) -Brauer diagram is a figure in the rectangle $R = I \times I$ consisting of

- (1) n distinguished points $\{1, \dots, n\}$ on the upper boundary $I \times \{1\}$ of R and n distinguished points $\{\bar{1}, \dots, \bar{n}\}$ on the lower boundary $I \times \{0\}$ of R .
- (2) n curves connecting the points $\{1, \dots, n, \bar{1}, \dots, \bar{n}\}$ in pairs, with each curve intersecting the boundary of R only at its two endpoints.


A $\mathbb{Z}_r$ -Brauer diagram is a Brauer diagram in which each curve (strand) is endowed with an orientation and labeled by an element of the group $\mathbb{Z}_r$. Two labelings are regarded as the same if the orientation of a strand is reversed and the group element associated to the strand is inverted. Note that the number of $\mathbb{Z}_r$ -Brauer diagrams is $r^n(2n-1)!!$.

Let A be an integral domain containing elements ϑ_j , for $0 \leq j \leq [r]$ with ϑ_0 invertible. The $\mathbb{Z}_r$ -Brauer algebra $D_{n,A,r}$ is defined as follows. As an A -module, $D_{n,A,r}$ is the free A -module with basis the set of $\mathbb{Z}_r$ -Brauer diagrams with n strands. The product of two $\mathbb{Z}_r$ -Brauer diagrams is defined to be a certain multiple of another $\mathbb{Z}_r$ -Brauer diagram, determined as follows.

Given two $\mathbb{Z}_r$ -Brauer diagrams a, b , first “stack” b over a (as for tangle diagrams and ordinary Brauer diagrams). In the resulting “tangle”, the strands are composed of one or more strands from a and b . Give each composite strand s an orientation, arbitrarily. Make the orientations of the components of s from a and b agree with the orientation of s , changing the signs of the $\mathbb{Z}_r$ -valued labels accordingly; the label of s is then the sum of the labels of its components from a and b . For each j ($0 \leq j \leq [r]$), let m_j be the number of closed loops with label equal to $\pm[j]$. Let c be the $\mathbb{Z}_r$ -Brauer diagram obtained by removing all the closed curves. Then $ab = (\prod_j \vartheta_j^{m_j}) c$. Extend the multiplication to a bilinear product on $D_{n,A,r}$. One can easily check that the multiplication is associative. Note that the subalgebra generated by $\mathbb{Z}_r$ -Brauer diagrams with only vertical strands is isomorphic to the group algebra of the wreath product $\mathbb{Z}_r \wr \mathfrak{S}_n$, so the $\mathbb{Z}_r$ -Brauer algebra can be considered as a sort of wreath product of $\mathbb{Z}_r$ with the ordinary Brauer algebra.

One has inclusion maps $\iota : D_{n-1,A,r} \rightarrow D_{n,A,r}$ defined on the level of $\mathbb{Z}_r$ -Brauer diagrams by adding a vertical strand on the right labeled by 0. Moreover, one has a conditional expectation $\varepsilon_n : D_{n,A,r} \rightarrow D_{n-1,A,r}$ defined as follows: First define a map cl_n , from $\mathbb{Z}_r$ -Brauer diagrams with n -strands to $\mathbb{Z}_r$ -Brauer diagrams with

$(n - 1)$ -strands, by joining the rightmost pair of vertices $\bar{n}, n$ of a diagram d by a new strand, with label 0:



Determine the $\mathbb{Z}_r$ -valued label of any concatenated strand in the resulting diagram by the same rule as in the multiplication of $\mathbb{Z}_r$ -Brauer diagrams. If the resulting diagram has a closed loop (which happens precisely if d contains a strand connecting n and $\bar{n}$ with some label $\pm[k]$) then remove the closed loop and multiply the resulting $\mathbb{Z}_r$ -Brauer diagram by ϑ_k . Finally, define $\varepsilon_n : D_{n,A,r} \rightarrow D_{n-1,A,r}$ by

$$\varepsilon_n(d) = \vartheta_0^{-1} \text{cl}_n(d).$$

One can check that ε_n is a conditional expectation. Define

$$\varepsilon = \varepsilon_1 \circ \cdots \circ \varepsilon_n : D_{n,A,r} \rightarrow D_{0,A,r} = A.$$

Then ε is a trace.

Lemma 5.1 ([25]). *For each j ($0 \leq j \leq \lfloor r \rfloor$), let ϑ_j be an indeterminant over $\mathbb{C}$. Let*

$$K = \mathbb{C}(\vartheta_0^{\pm 1}, \vartheta_1, \dots, \vartheta_{\lfloor r \rfloor}).$$

The trace ε on $D_{n,K,r}$ is non-degenerate.

Given this, it is fairly straightforward to apply Wenzl's method from [29, 30] to the $\mathbb{Z}_r$ -Brauer algebra. (In fact, this has been done more generally in [25], in the context of G -Brauer algebras, where G is a finite abelian group.) The result of this analysis is that the $\mathbb{Z}_r$ -Brauer algebra $D_n = D_{n,K,r}$ is split semisimple with simple modules labelled by Γ_n (i.e., the set of r -tuples of Young diagrams of total size $\leq n$ and congruent to $n \bmod 2$) and the branching diagram for the sequence $D_0 \subseteq D_1 \subseteq \cdots \subseteq D_n$ is exactly the same as that obtained for the sequence of cyclotomic BMW algebras in Theorem 4.8. Thus we have:

Theorem 5.2. *Let F be as in Theorem 4.8. The dimension of $W_{n,F,r}$ is $r^n(2n - 1)!!$.*

Proof. $\dim_F(W_{n,F,r}) = \sum_{\lambda \in \Gamma_n} |\mathcal{T}(n, \lambda)|^2 = \dim_K(D_{n,K,r}) = r^n(2n - 1)!!$. $\square$

6. FREENESS AND ISOMORPHISM WITH CYCLOTOMIC KAUFFMAN TANGLE ALGEBRAS.

We show in [10], Proposition 3.6, that every cyclotomic BMW algebra $W_{n,S,r}(u_1, \dots, u_r)$ is spanned over its ground ring S by a set $\mathbb{A}'_r$ of cardinality $r^n(2n-1)!!$. Combining this result with Theorems 4.8 and 5.2, we obtain the following theorem (which has been proved independently by Wilcox and Yu.)

Theorem 6.1 (Goodman and Hauschild–Mosley, Wilcox and Yu [33, 32]). *Let S be an admissible integral domain. Then $W_{n,S,r} \cong KT_{n,S,r}$, and $W_{n,S,r}$ is a free S -module of rank $r^n(2n-1)!!$.*

Proof. As noted above, $KT_{n,S,r}$ has a spanning set $\mathbb{A}'_r$ of cardinality $r^n(2n-1)!!$. To prove the theorem, it suffices to show that $\varphi(\mathbb{A}'_r) = \mathbb{B}'_r$ is linearly independent in $KT_{n,S,r}$. When $S = F$, the field of fractions of the universal admissible integral domain $\overline{R}_+$, this follows from Theorem 4.8 and Theorem 5.2, since $\mathbb{B}'_r$ is a spanning set whose cardinality equals the dimension of $KT_{n,F,r}$. The map $x \mapsto x \otimes 1$ from $KT_{n,\overline{R}_+,r}$ to $KT_{n,\overline{R}_+,r} \otimes_{\overline{R}_+} F \cong KT_{n,F,r}$ is $\overline{R}_+$ -linear and maps $\mathbb{B}'_r$ to a linearly independent set in $KT_{n,F,r}$; hence $\mathbb{B}'_r$ is linearly independent in $KT_{n,\overline{R}_+,r}$. Finally, since any admissible integral domain S is a quotient of $\overline{R}_+$, and $KT_{n,\overline{R}_+,r}$ is a free $\overline{R}_+$ -module with basis $\mathbb{B}'_r$, it follows that $KT_{n,S,r} \cong KT_{n,\overline{R}_+,r} \otimes_{\overline{R}_+} S$ is a free S -module with basis $\mathbb{B}'_r$. $\square$

Corollary 6.2. *Suppose S is an admissible integral domain.*

- (1) *There is a trace-preserving conditional expectation $\varepsilon_n : W_{n+1,S,r} \rightarrow W_{n,S,r}$.*
- (2) *The natural map $\iota : W_{n,S,r} \rightarrow W_{n+1,S,r}$ is injective.*

REFERENCES

1. Susumu Ariki, *Representations of quantum algebras and combinatorics of Young tableaux*, University Lecture Series, vol. 26, American Mathematical Society, Providence, RI, 2002, Translated from the 2000 Japanese edition and revised by the author. MR MR1911030 (2004b:17022)
2. Susumu Ariki and Kazuhiko Koike, *A Hecke algebra of $(\mathbb{Z}/r\mathbb{Z}) \wr \mathfrak{S}_n$ and construction of its irreducible representations*, Adv. Math. **106** (1994), no. 2, 216–243. MR MR1279219 (95h:20006)
3. Susumu Ariki, Andrew Mathas, and Hebing Rui, *Cyclotomic Nazarov–Wenzl algebras*, Nagoya Math. J. **182** (2006), 47–134. MR MR2235339
4. Michael Artin, *Algebra*, Prentice Hall Inc., Englewood Cliffs, NJ, 1991. MR MR1129886 (92g:00001)
5. Anna Beliakova and Christian Blanchet, *Skein construction of idempotents in Birman–Murakami–Wenzl algebras*, Math. Ann. **321** (2001), no. 2, 347–373. MR MR1866492 (2002h:57018)

6. Joan S. Birman and Hans Wenzl, *Braids, link polynomials and a new algebra*, Trans. Amer. Math. Soc. **313** (1989), no. 1, 249–273. MR 90g:57004
7. Frederick M. Goodman, *Cellularity of cyclotomic Birman–Wenzl–Murakami algebras*, preprint (2008), arXiv:0801.0306.
8. Frederick M. Goodman, Pierre de la Harpe, and Vaughan F. R. Jones, *Coxeter graphs and towers of algebras*, Mathematical Sciences Research Institute Publications, vol. 14, Springer-Verlag, New York, 1989. MR MR999799 (91c:46082)
9. Frederick M. Goodman and Holly Hauschild, *Affine Birman–Wenzl–Murakami algebras and tangles in the solid torus*, Fund. Math. **190** (2006), 77–137. MR MR2232856
10. Frederick M. Goodman and Holly Hauschild Mosley, *Cyclotomic Birman–Wenzl–Murakami algebras I: Freeness and realization as tangle algebras*, preprint (2006, Revised 2007), arXiv:math/0612064.
11. J. J. Graham and G. I. Lehrer, *Cellular algebras*, Invent. Math. **123** (1996), no. 1, 1–34. MR MR1376244 (97h:20016)
12. Reinhard Häring-Oldenburg, *Cyclotomic Birman–Murakami–Wenzl algebras*, J. Pure Appl. Algebra **161** (2001), no. 1–2, 113–144. MR MR1834081 (2002c:20055)
13. V. F. R. Jones, *Index for subfactors*, Invent. Math. **72** (1983), no. 1, 1–25. MR MR696688 (84d:46097)
14. Louis H. Kauffman, *An invariant of regular isotopy*, Trans. Amer. Math. Soc. **318** (1990), no. 2, 417–471. MR MR958895 (90g:57007)
15. C. Krattenthaler, *Advanced determinant calculus*, Sémin. Lothar. Combin. **42** (1999), Art. B42q, 67 pp. (electronic), The Andrews Festschrift (Maratea, 1998). MR MR1701596 (2002i:05013)
16. Sofia Lambropoulou, *A study of braids in 3-manifolds*, Ph.D. Thesis, Warwick (1993).
17. ———, *Solid torus links and Hecke algebras of $\mathcal{B}$ -type*, Proceedings of the Conference on Quantum Topology (Manhattan, KS, 1993), World Sci. Publ., River Edge, NJ, 1994, pp. 225–245. MR MR1309934 (96a:57020)
18. Robert Leduc and Arun Ram, *A ribbon Hopf algebra approach to the irreducible representations of centralizer algebras: the Brauer, Birman–Wenzl, and type A Iwahori–Hecke algebras*, Adv. Math. **125** (1997), no. 1, 1–94. MR MR1427801 (98c:20015)
19. Andrew Mathas, *Seminormal forms and Gram determinants for cellular algebras*, to appear in J. Reine Angew. Math., with an appendix by Marcos Soriano.
20. Hugh Morton and Paweł Traczyk, *Knots and algebras*, Contribuciones Matemáticas en homenaje al profesor D. Antonio Plans Sanz de Bremond (E. Martín-Peindador and A. Rodez Usan, eds.), University of Zaragoza, Zaragoza, 1990, pp. 201–220.
21. Hugh Morton and Antony Wassermann, *A basis for the Birman–Wenzl algebra*, Unpublished manuscript (1989, revised 2000), 1–29.
22. Jun Murakami, *The Kauffman polynomial of links and representation theory*, Osaka J. Math. **24** (1987), no. 4, 745–758. MR MR927059 (89c:57007)
23. Maxim Nazarov, *Young’s orthogonal form for Brauer’s centralizer algebra*, J. Algebra **182** (1996), no. 3, 664–693. MR MR1398116 (97m:20057)
24. Rosa Orellana and Arun Ram, *Affine braids, Markov traces and the category $\mathcal{O}$* , preprint, 2004, arXiv:math.RT/0401317.
25. M. Parvathi and D. Savithri, *Representations of G -Brauer algebras*, Southeast Asian Bull. Math. **26** (2002), no. 3, 453–468. MR MR2047837 (2005b:16056)
26. Hebing Rui and Jie Xu, *The representations of cyclotomic BMW algebras*, preprint (2008), arXiv:0801.0465.
27. Hebing Rui and Weihua Yu, *On the semi-simplicity of the cyclotomic Brauer algebras*, J. Algebra **277** (2004), no. 1, 187–221. MR MR2059627 (2005b:16057)

- 28. Vladimir Turaev, *The Conway and Kauffman modules of a solid torus*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **167** (1988), no. Issled. Topol. 6, 79–89, 190. MR MR964255 (90f:57012)
- 29. Hans Wenzl, *On the structure of Brauer’s centralizer algebras*, Ann. of Math. (2) **128** (1988), no. 1, 173–193. MR MR951511 (89h:20059)
- 30. ———, *Quantum groups and subfactors of type B, C, and D*, Comm. Math. Phys. **133** (1990), no. 2, 383–432. MR MR1090432 (92k:17032)
- 31. Stewart Wilcox and Shona Yu, *The cyclotomic BMW algebra associated with the two string type B braid group*, preprint (2006), arXiv:math/0611518.
- 32. ———, in preparation (2008).
- 33. Shona Yu, *The cyclotomic Birman–Murakami–Wenzl algebras*, Ph.D. Thesis, University of Sydney (2007).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, IOWA CITY, IOWA
E-mail address: `goodman@math.uiowa.edu`

DEPARTMENT OF MATHEMATICS, GRINNELL COLLEGE, GRINNELL, IOWA
E-mail address: `HAUSCHIL@GRINNELL.EDU`